

PRODUCT LITERATURE

GUAICOL SYRUP

Each 5ml contains

Guaiphenesin USP 100mg

Preservatives:

Methyl Paraben BP 0.1% w/v

Propyl Paraben BP 0.01% w/v

Product description

Red coloured clear syrup with fruity odour and sweet taste.

Pharmacodynamic properties

Guaifenesin reduces the viscosity of tenacious sputum and is used as an expectorant

Pharmacokinetics properties

Guaifenesin is readily absorbed from the gastrointestinal tract. It is metabolised and excreted in the urine.

Indication

As an expectorant for the symptomatic relief of cough due to colds and minor upper respiratory infections.

Recommended dose

To be given 3 to 4 times a day.

Adult : 10 – 20 ml (2-4 tsp)

Children (6-12 years) : 5 - 10 ml (1-2 tsp)

(2-6 years) : 2.5 – 5 ml (½-1 tsp)

Route of administration

Oral

Contradictions

Hypersensitivity to the active substance or any of the excipients.

Warning and Precautions

Do not exceed the stated dose.

Ask a doctor before use if you suffer from a chronic cough, if you have asthma or are suffering from an acute asthma attack.

Stop use and ask a healthcare professional if your cough lasts for more than 5 days, comes back, or is accompanied by a fever, rash or persistent headache.

Do not take with a cough suppressant.

Interaction with other medicaments

There are no clinically significant interactions.

If urine is collected within 24 hours of a dose of guaifenesin a metabolite of guaifenesin may cause a colour interference with laboratory determinations of urinary 5-hydroxyindoleacetic acid (5-HIAA) and vanillylmandelic acid (VMA).

Pregnancy and Lactation

The safety of this product during pregnancy and lactation has not been established. However guaiphenesin is not considered to constitute a hazard during pregnancy since there is no evidence of an association with foetal malformations. There is no data available on use of guaiphenesin during breast feeding.

Side effects

The following side effects may be associated with the use of guaifenesin:

Gastrointestinal disorders: nausea, vomiting/gastrointestinal discomfort.

Immune system disorder: hypersensitivity reactions, including anaphylaxis.

Storage Condition

Store below 30°C, in a dry place, protected from direct light.

Keep away from reach of children.

Jauhkan daripada kanak- kanak

SHAKE WELL BEFORE USE

Shelf life

2 years from the manufacturing date

Name and Address of Product Registration Holder/

Distributor:

ZONTRON PHARMACEUTICALS SDN.BHD. (445695-T)

Lot 10 & 11, PERDA Industrial Park,

Lorong IKS Simpang Ampat B,

14100 Simpang Ampat, S.P.S.,

Pulau Pinang, Malaysia

Name and Address of Manufacturer:

TERAPUTICS SDN. BHD. (590500-W)

Lot 10 & 11, PERDA Industrial Park,

Lorong IKS Simpang Ampat B,

14100 Simpang Ampat, S.P.S.,

Pulau Pinang, Malaysia

Malaysian Drug Registration No. MAL19920118XC

Date of Revision: 08/10/2019