



60/12623685/0217

Directions for use
Read carefully!

B. Braun 0.45% Sodium Chloride and 10% Glucose Intravenous Infusion B.P.

Composition

Each 100 ml contains:
Sodium Chloride 0.45 g
Glucose Monohydrate for Parenteral Use 11.0 g
Water for Injections to 100 ml
Electrolytes: mmol/l (mEq/l)
Na⁺ 77
Cl⁻ 77
Caloric value: 1700 kJ/l $\triangle$ 400 kcal/l
Osmolarity: 709 mOsm/l

Indications

Hypertonic (especially hypernatremia) dehydration
Sodium and chloride depletion
Caloric supply
Vehicle solution for supplementary medication

Contraindications

Hyperhydration, oedema
Hypernatremia
Hypokalaemia

Dosage

According to individual requirements
approximately 1000 ml/day.
Drop rate: up to 120-180 drops/min
corresponding to 360-540 ml/h.

Route of administration I.V.

Precautions

Use with caution in patients with hypertension and diabetes mellitus. Blood glucose, serum electrolytes and water balance should be monitored regularly.

The compatibility of any additives to this solution should be checked before use.

Side effects

Hyperglycaemia and glucosuria may occur if glucose tolerance is impaired. These can be avoided by reducing the dosage and infusion rate or by administering insulin.

Symptoms and treatment for overdosage

If hyperglycaemia or glucosuria occurs, the dosage and infusion rate should be reduced, or insulin may be administered.

Shelf life

The product must not be used beyond the expiry date stated on the label.

Storage

Do not store above 25°C.

Presentation

500 ml, 1000 ml plastic container.

Approval for Printing

B|BRAUN Melsungen AG

Approved for Printing ☐

Approved for Printing
when corrected ☐

New draft required ☐

Date

Signature

Name in capital letters

schwarz

Dokument = 148 x 210 mm (DIN A5)
1 Seite

MY__60
GIF (EP)
0.45% Sodium Chloride and 10% Glucose
60/12623685/0217
Standort Penang

Lätus



8771

Font size: 9.25 pt.

G 110921

B|BRAUN

Product registration holder
and manufactured by:

B. Braun Medical
Industries Sdn. Bhd.
11900 Bayan Lepas,
Penang, Malaysia.

