



1059/12623710/0717

Directions for use

Read carefully!

Composition

Each 100 ml contains:

Sodium Chloride	0.20 g
Glucose Monohydrate for Parenteral Use	5.50 g
Water for Injections to	100 ml

Electrolytes:	mmol/l (mEq/l)
Na ⁺	34
Cl ⁻	34
Caloric value:	850 kJ/l $\triangleq$ 200 kcal/l
Osmolarity:	346 mOsm/l

Characteristics

B. Braun 0.20% Sodium Chloride and 5% Glucose Intravenous Infusion B.P. is a light hypertonic solution containing 34 mmol each of sodium and chloride per litre and supplies physiologically free water.

Mechanisms of action

Glucose is absorbed from the gastrointestinal tract, and oxidised as a source of energy, or stored in the liver as glycogen. It is the only energy substrate which is directly, instantly and universally utilized by the body. Glucose is vital for the myocardium, brain and nerves.

Sodium Chloride is the principal salt involved in maintaining the osmotic tension of the blood and tissue. Changes in sodium and chloride levels change this osmotic tension and hence influence the movement of fluids and diffusion of salts in cellular tissue.

Indications

Dehydration,
Sodium and chloride depletion,
Caloric supply,
Vehicle solution for supplementary medication.

Dosage

According to individual requirements approximately 1000 ml/day for adult patients.
Drop rate: up to 120-180 drops/min corresponding to 360-540 ml/h

B. Braun 0.20% Sodium Chloride and 5% Glucose Intravenous Infusion B.P.

Contraindications

Hyperhydration, oedema
Hypernatremia
Hypokalaemia

Route of administration I.V.

Precautions

Use with caution in patients with hypertension and diabetes mellitus. Blood glucose, serum electrolytes and water balance should be monitored regularly.

Side effects

Hyperglycaemia and glucosuria may occur if glucose tolerance is impaired. These can be avoided by reducing the dosage and infusion rate or by administering insulin.

Symptoms and treatment for overdosage

If hyperglycaemia or glucosuria occurs, the dosage and infusion rate should be reduced, or insulin may be administered.

Usage during pregnancy

No adverse reaction has been reported with sodium chloride content. However, the glucose content indicated rapid infusions of solutions with 25g or more glucose lead to foetal acidosis and neonatal hyperinsulinaemia hypoglycaemia and jaundice. It is recommended to limit infusion to not more than 6g glucose per hour just before delivery, until a safe rate of administration is established.

Shelf life

The product must not be used beyond the expiry date stated on the label.

Storage

The product should not be stored above the temperature stated on the label.

Presentation

500 ml plastic container.
Not all presentation is available locally.

Approval for Printing

B. BRAUN Melsungen AG

Approved for Printing ☐

Approved for Printing when corrected ☐

New draft required ☐

Date _____ Signature _____

Name in capital letters

schwarz

Format = 148 x 210 mm (DIN A5)

1 Seite

Lätus



5403

MY__1059

1059/12623710/0717

GIF (EP)

Standort Penang

Font size: 8 pt.

G 170200

Product Registration Holder and
Manufactured by:

B. Braun Medical Industries Sdn. Bhd.
11900 Bayan Lepas, Penang,
Malaysia.

B. BRAUN

