

HOVID-GRISEOFULVIN TABLET 500mg

VIGRI08-var (MY)

DESCRIPTION

Round, white film-coated tablet, bevel-edged, shallow convex faces, "HD" embossed and scored on the same face.

COMPOSITION

Each tablet contains:
Griseofulvin 500 mg.

PHARMACODYNAMICS

Griseofulvin inhibits fungal cell mitosis by causing disruption of the mitotic spindle structure, thereby arresting the metaphase of cell division. It is deposited in the keratin precursor cells of skin, hair, and nails rendering the keratin resistant to fungal invasion. As the infected keratin is shed, it is replaced with healthy tissue.

PHARMACOKINETICS**Absorption**

Irregularly and incompletely absorbed after oral administration; absorption of griseofulvin is particle-size dependent; its absorption is increased when administered with fatty meals or milk and increased when administered concomitantly with phenobarbitone.

Blood Concentration

After an oral daily dose of 0.5 to 1.0g, peak plasma concentrations of 1 to 3mg/ml are attained; after single oral doses of 0.25 to 3g. Peak plasma concentrations of 0.3 to 1.7mg/ml are attained in 4 hours.

Half-life

Plasma half-life for the alpha-phase, about 1 hour and for the beta-phase, 9 to 22 hours.

Distribution

Localised in the skin, nails, and hair roots; secreted in the sweat and crosses the placenta.

Metabolic Reactions

O-Demethylation at the 4-and 6-positions and glucuronic acid conjugation of the demethylated-metabolites.

Excretion

After an oral dose, about 35% is excreted in the urine as the 6-demethylated metabolite in 8 days; after an intravenous dose about 65% is excreted in the urine as the same metabolite in 72 hours; large amounts of an oral dose are eliminated unchanged in the faeces but less than 1% is excreted unchanged in the urine; the urinary excretion of 6-demethylgriseofulvin appears to be pH-dependent; small amounts of the 4-demethylated metabolite are excreted in the urine.

INDICATIONS

For treatment of dermatophytic infections of the skin, nails and hair caused by species of Epidermophyton, Microsporum and Trichophyton.

CONTRAINDICATIONS

- Not recommended for patients with porphyria and liver failure.
- Griseofulvin is contraindicated in pregnancy.

WARNING AND PRECAUTIONS

- Caution in patients hypersensitive to penicillins and those with lupus erythematosus.
- Safety for use in pregnancy and lactation has not been established.
- Griseofulvin may damage sperm cells; male patients should not father children during treatment or within six months of completion of treatment.
- Some case reports of human foetal abnormalities have been observed. Therefore, women should not become pregnant during or within one month of treatment with griseofulvin.
- Severe cutaneous adverse reactions (e.g. Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms, acute generalised exanthematous pustulosis) and erythema multiforme have been reported with griseofulvin use. These reactions may be serious and may result in hospitalisation or death. If severe skin reactions occur, griseofulvin should be discontinued.

PREGNANCY AND LACTATION

Safety for use in pregnancy and lactation has not been established.

DRUG INTERACTIONS

- Concurrent use with alcohol may result in tachycardia and flushing.
- Co-administration of barbiturates may decrease effects of griseofulvin.
- Effects of coumarin-type anticoagulants may be decreased when used concurrently with griseofulvin.
- The effectiveness of oral contraceptives may be reduced during treatment with griseofulvin. It may be advisable to use alternative or additional method of contraception during therapy and for one month after stopping of griseofulvin.

MAIN SIDE/ADVERSE EFFECTS

Headache, hypersensitivity reaction, dryness of the mouth and gastro-intestinal disturbances.

Skin and subcutaneous tissue disorders

Frequency 'not known': Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms, acute generalised exanthematous pustulosis, erythema multiforme.

OVERDOSAGE**Clinical features**

Urticaria, nausea, diarrhoea, headaches, confusion, dizziness, photosensitivity, renal damage, porphyria and temporary leukopenia.

Treat gastrointestinal distress, if any, by giving milk every 3 hours alternating with bismuth subcarbonate 5 g every 3 hours.

DOSAGE AND ADMINISTRATIONAdults:

Oral, 500 mg per day as a single dose or in divided doses, or as directed.

Children:

Oral, 10 mg/kg of body weight per day as a single dose or in divided doses, or as directed.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage: Store below 30°C. Protect from moisture.

Presentation/Packing:

Film-coated tablet 500 mg x 100's, 500's, 1000's,
Blisters of 10 x 10's, Blisters of 2 x 10's.

(Not all presentation available locally)

Product Registration Holder (Malaysia) /

Product Owner: HOVID Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),

30010 Ipoh, Perak, Malaysia.

Manufactured by: HOVID Bhd.

Lot 56442, 7 1/2 Miles, Jalan Ipoh/Chemor,

31200 Chemor, Perak, Malaysia.

Revision date: October 2023



20109



20109