

BUTYLIN

VIBUT01-var

DESCRIPTION

Colourless, transparent liquid with cherry flavour.

COMPOSITION

Each 5 ml contains: Terbutaline Sulphate 1.5 mg

ACTIONS AND PHARMACOLOGY

Terbutaline is a direct-acting sympathomimetic agent with actions on beta-2-adrenergic receptors in the lungs to relax bronchial smooth muscle, thereby relieving bronchospasm and reducing airway resistance.

Oral absorption is incomplete, with fairly extensive first-pass hepatic metabolism. Renal excretion is via the kidneys.

INDICATIONS

For the management of asthma, chronic bronchitis, emphysema and other broncho-pulmonary disorders involving bronchospasm.

CONTRAINDICATIONS

- Not recommended for hypersensitive patients.
- Avoid concurrent therapy with non-selective beta-adrenoreceptor blockers.
- Oral terbutaline sulfate is contraindicated for the treatment of acute or maintenance tocolysis.

WARNING AND PRECAUTIONS

- Caution in patients with cardiac disorders, especially those associated with arrhythmias, diabetes mellitus, hypertension, hyperthyroidism and history of seizures.
- Check with physician immediately if difficulty in breathing persists after using this medication or if condition becomes worse.
- There are no adequate and well-controlled studies of terbutaline sulfate in pregnant women. Published animal studies show that rat offspring exhibit alterations in behaviour and brain development, including decreased cellular proliferation and differentiation when dams were treated subcutaneously with terbutaline during the late stage of pregnancy and lactation period. Terbutaline exposures in rat dams were approximately 6.5 times the common human dose in adults of 15 mg/day, on a mg/m² basis.
- Tocolysis: Serious adverse reactions including death have been reported after administration of terbutaline/ salbutamol to women in labor. In the mother, these include increased heart rate, transient hyperglycaemia, hypokalaemia, cardiac arrhythmias, pulmonary oedema and myocardial ischaemia. Increased fetal heart rate and neonatal hypoglycaemia may occur as a result of maternal administration.

MAIN SIDE/ADVERSE EFFECTS

Nervousness, restlessness, muscle tremors, headaches, nausea or vomiting.

DRUG INTERACTIONS

Concomitant administration of other sympathomimetics may enhance effects of either medications.

OVERDOSAGE

Clinical features:

Tremor, palpitations, tachycardia, peripheral vasodilatation, hypokalaemia, hyperglycaemia, lactic acidosis.

Treatment for overdosage:

Emesis or gastric lavage, if appropriate. Tachycardia and myocardial ischaemia may be diminished by administering atenolol 2.5-10 mg or propranolol 1 mg/min slowly (caution in asthmatic patients).

DOSAGE AND ADMINISTRATION

Adults

Oral, 10 ml to 15 ml three times a day, up to 50 ml per day.

Children

Below 12 years : As directed by physician.
12 -15 years : Oral, 10 ml three times a day.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage:

Store below 30°C.

Presentation/Packing:

Syrup x 60 ml, 100 ml, 120 ml, 3.6 Litres.
(Not all presentation available locally)

Product Registration Holder (Malaysia) /

Manufactured by: HOVID Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia

Revision date: May 2025