

ORALON Dental Paste

NAME AND STRENGTH OF ACTIVE SUBSTANCE:

ORALON dental paste is Triamcinolone Acetonide 0.1% w/w in emollient dental paste.

Each gram contains:

Triamcinolone Acetonide.....1mg

This product contains gelatin of non-porcine origin.

PRODUCT DESCRIPTION:

A light brown paste in printed tubes with white caps.

PHARMACODYNAMICS:

Triamcinolone acetonide is used in dermatology for its anti-inflammatory, antipruritic and vasoconstrictive action on the skin/mucous membranes.

PHARMACOKINETICS:

Topically applied triamcinolone acetonide is usually metabolised in the skin. There is some systemic absorption through the oral mucosa. Once the drug is absorbed into the systemic circulation, it is bound to plasma proteins. It is mainly metabolised in the liver.

INDICATION:

ORALON Dental Paste is indicated for adjunctive treatment and temporary relief of symptoms associated with non-herpetic, oral inflammatory and ulcerative lesions such as: recurrent ulcerative stomatitis, erosive lichen planus, denture stomatitis, traumatic lesions including denture sore spots, desquamative gingivitis and stomatitis, and aphthous stomatitis.

RECOMMENDED DOSAGE:

Usual dose:

Apply a small dab of the ORALON Dental Paste (about ½ cm) to the lesion, two or three times a day, after meals, and at bed time.

Applying the paste with a cotton applicator will help to eliminate any possible absorption from contact with skin.

The paste should be pressed, not rubbed, on the lesion. Rubbing the paste on the lesion will result in a granular, gritty sensation and cause it to crumble. A smooth, slippery film forms after application.

ROUTE OF ADMINISTRATION:

To be used topically, to the oral mucous membranes.

CONTRAINDICATIONS:

- a) The use of topical corticosteroids, is contraindicated in herpetic lesions of known viral origin such as herpes labialis, intraoral lesions such as primary herpetic gingival stomatitis and herpanginas.
- b) Known hypersensitivity to any ingredient of the preparation.

WARNINGS AND PRECAUTIONS:

- i) In general corticosteroids should not be used in the presence of infection.
- ii) Patients with tuberculosis, peptic ulcer, or diabetes mellitus should not be treated with corticosteroid preparations, including the dental paste.
- iii) It should be borne in mind that the normal defensive responses of the oral tissues are depressed in patients receiving topical corticosteroid therapy.
- iv) If significant regeneration or repair of oral tissues has not occurred in 7 days, additional investigations into the etiology of oral lesion is advised.

INTERACTIONS WITH OTHER MEDICAMENTS:

Unknown

STATEMENT ON USAGE DURING PREGNANCY:

Should not be used in women of child-bearing potential and particularly during early pregnancy unless the potential benefits outweighed the possible hazards.

ADVERSE EFFECTS/UNDESIRABLE EFFECTS:

- a) Signs of infection such as pain, redness or pus-containing blister.
- b) Signs of irritation such as burning, itching, blistering, or peeling not present before therapy.

OVERDOSAGE AND TREATMENT:

Symptoms: Signs of irritation such as burning, itching, blistering or peeling.

Treatment: Withdrawal of the drug is usually all that is necessary.

DOSAGE FORMS AND PACKAGING AVAILABLE:

Packing size: Collapsible aluminium tube of 5g.

The tube should be tightly closed after each administration.

Keep out of reach of children. / Jauhi dari kanak-kanak.

NAME AND ADDRESS OF MANUFACTURER / PRODUCT REGISTRATION HOLDER:

MALAYSIAN PHARMACEUTICAL INDUSTRIES SDN BHD (101323-U)

Plot 14, Lebuhraya Kampung Jawa, 11900 Bayan Lepas,

Pulau Pinang, Malaysia.

DATE OF REVISION:

22/05/2017