

(L)210mm

pharmaniaga®



Terbutaline

Tablet

DESCRIPTION

White to off white, round, biconvex tablet with 'RMB' breakline on one side and diamond shape marking on the other.

COMPOSITION

Each tablet contains Terbutaline sulphate 2.5 mg.

ACTIONS

Terbutaline sulphate is a direct-acting sympathomimetic agent with actions and uses similar to those of salbutamol.

Salbutamol is a direct-acting sympathomimetic agent with a predominantly beta₂-adrenergic activity and a selective action on beta₂ receptors.

Terbutaline is incompletely absorbed from the gastrointestinal tract and is also subject to fairly extensive first-pass metabolism by sulphate (and some glucuronide) conjugation in the liver and possibly the gut wall. It is accordingly excreted in the urine partly as the inactive conjugates and partly as unchanged terbutaline, the ratio depending upon whether it was given by mouth or parenterally.

INDICATIONS

Indicated for the relief and prevention of bronchial asthma.

ADVERSE REACTIONS

Sympathomimetics may produce a wide range of adverse effects, most of which mimic the results of

excessive stimulation of the sympathetic nervous system. These effects are mediated via the various type of adrenergic receptor and the adverse effects of an individual drug depend to some extent upon its relative agonist activity on these different type of receptors at a given dose.

Central effects of sympathomimetic agents include fear, anxiety, restlessness, tremor, insomnia, confusion, irritability, weakness and psychotic states. Appetite may be reduced, and nausea and vomiting may occur.

Effects on the cardiovascular system are complex. Stimulation of alpha-adrenergic receptor produces vasoconstriction, sometimes sufficiently severe to produce gangrene when infiltrated into the digits with resultant hypertension; the rise in blood pressure may produce cerebral haemorrhage and pulmonary oedema. There may also be a reflex bradycardia, but stimulation of beta₂-adrenergic receptors of the heart may produce tachycardia and cardiac arrhythmias, anginal pain, palpitation and cardiac arrest; hypotension with dizziness and fainting and flushing may occur. An increased incidence of sudden death, perhaps as a result of the induction of ventricular arrhythmias, has been associated with the excessive use of sympathomimetic agent in aerosol form; although the association has been questioned by some authorities it is important to avoid excessive doses.

Other effects that may occur with sympathomimetic agents include difficulty in micturition and urinary retention, dyspnoea, altered metabolism including disturbance of glucose metabolism, sweating and hypersalivation. Headache is also common.

CONTRAINDICATIONS

In patients with known hypersensitivity to the drug.

Effects On Ability To Drive and Use Machine

Terbutaline has no or negligible influence on the ability to drive and use machines.

PRECAUTIONS

Tocolysis: Serious adverse reactions including death have been reported after administration of terbutaline/salbutamol to women in labour. In the mother, these include increased heart rate, transient hyperglycaemia, hypokalaemia, cardiac arrhythmias, pulmonary oedema and myocardial ischaemia. Increased fetal heart rate and neonatal hypoglycaemia may occur as a result of maternal administration.

As maternal pulmonary oedema and myocardial ischaemia have been reported during or following premature labour in patients receiving beta₂-agonists, careful attention should be given to fluid balance and cardio-respiratory function, including ECG monitoring. If signs of pulmonary oedema and myocardial ischaemia develop, discontinuation of treatment should be considered.

Due to the risk of pulmonary oedema and myocardial ischaemia that has been observed during the use of beta₂-agonists in the treatment of premature labour, before these products are given to any patients with known heart disease, an adequate assessment of the patients cardiovascular status should be made by a physician experienced in cardiology.

Sympathomimetic agents should be used with caution in patients who may be hypersusceptible to their effect, particularly those with hyperthyroidism. Great care is also needed in patients with cardiovascular disease: arrhythmia or tachycardia; occlusive vascular disorder, including arteriosclerosis, hypertension or aneurysms. Anginal pain may be precipitated in patients with angina pectoris. Care is also required when sympathomimetics are given to patients with diabetes mellitus or closed-angle glaucoma.

Tolerance may be developed in asthmatic patients given sympathomimetics for their bronchodilator effects; if tolerance occurs and the patients condition worsens, alternative or additional therapy should be instituted. The dose of the sympathomimetics should not be increased in such cases. Sympathomimetics should be avoided or used with caution in patients undergoing anaesthesia with cyclopropane, halothane or other halogenated anaesthetics, as they may induce ventricular fibrillation. An increased risk of arrhythmias may also occur if sympathomimetic agents are given to patients receiving cardiac glycosides, quinidine or tricyclic antidepressants.

DOSAGE AND ADMINISTRATION

Adults:

Initially 2.5 mg three times daily increased to 5 mg three times daily as necessary.

Children over 7 years:

2.5 mg two or three times daily or 75 mcg per kg body-weight three times daily.

OVERDOSAGE

Symptoms include tremor and tachycardia which could be treated with a beta-adrenoceptor antagonist, such as propranolol.

ROUTE OF ADMINISTRATION

Oral.

STORAGE CONDITIONS

Store below 30°C.

Protect from light.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PRESENTATION

In blisters of 500 tablets.

Date of revision: 13th January 2021

PRODUCT REGISTRATION HOLDER/ MANUFACTURER:

PHARMANIAGA MANUFACTURING BERHAD
(198001006232)

No. 11A, Jalan P/1, Kawasan Perusahaan Bangi,
43650 Bandar Baru Bangi, Selangor Darul Ehsan,
Malaysia

PRP 0061.15 130121

attn

customer **Pharmaniaga Manufacturing Berhad**date **10.05.2021****1** coloursprocess
colours:**Black****Please Chop & Sign For Approval**

size (L)210 x (W)200mm

description **Terbutaline Insert****(PRP 0061.15 130121)**material
finishing**60g Simili****FocusPrint SDN BHD**t: 603-8766 6030 f: 603-8766 6033 (Factory)
website: www.focusprint.com.my

artwork prepared by: cynthia yap

email: graphic@focusprint.info (Graphic Dept)

date:

IMPORTANT ! Please note that this colour print is for visual purpose only. Output colour might differ in actual production.

ARTWORK LOG

Version no.	Date	Reason for Change
01	19.09.17	- Amend wording & PRP no.
02	29.09.17	- Amend wording.
03	09.10.17	- Amend wording.
04	24.10.17	- Amend wording at Effects On Ability To Drive and Use Machine
05	13.01.2021	- Amend wording.
06	19.02.2021	- Add BE Logo.

Version no.	Date	Reason for Change
07	10.05.2021	- Amend BE icon location.

(W)200mm