

PROCELL E 400

Description

Oval, transparent soft gelatin capsule filled with yellow oil.

Composition

d-alpha tocopherol 400 i.u

Action and Pharmacology

Vitamin E is considered as an essential nutritional element. As an antioxidant, it prevents the actions of peroxidase on unsaturated bonds in cell membranes and protects red blood cells against haemolysis.

Vitamin E is well absorbed from the gastrointestinal tract. It is metabolised in the liver and is excreted in the bile and urine.

Indications

Vitamin E is indicated for the treatment of Vitamin E deficiency.

Precautions

Safety for use in pregnancy and lactation has not been established.

Main Side/Adverse Effects

Vitamin E is usually well tolerated. Large doses greater than 400 to 800 units per day for prolonged periods may cause.

- Flu-like symptoms
- Pain in lower legs (thrombophlebitis)
- Stomach pain, tiredness, weakness
- Blurred vision
- Breast enlargement in males and females
- Diarrhoea, dizziness, headache and nausea

Note:

Very high doses of vitamin E (greater than 800 units per day for prolonged periods) have also been associated with increase bleeding tendencies in Vitamin K deficient patients, altered metabolism of hormones (thyroids, pituitary and adrenal), altered immunity, impaired sexual function and may increase the risk of thrombophlebitis or thromboembolism in susceptible patients (although this is controversial).

Drug Interactions

- Concurrent use with iron supplements should be avoided since Vitamin E impairs the hematologic response in patients with iron deficiency anaemia.
- Vitamin E should not be used concurrently with Vitamin A since large doses of Vitamin E deplete Vitamin A stores.
- Concurrent use with Vitamin K is not recommended since Vitamin E antagonises the effect of Vitamin K.

Overdosage

Clinical Features : Diarrhoea, abdominal pain and other gastrointestinal disturbances, flu-like symptoms, pain in lower legs, fatigue and weakness.

Treatment : Discontinue the medication immediately. Seek advice from your doctor

Dosage and Administration

Adults : Oral, one capsule daily

Note : The information given here is limited. For further information consult your doctor or pharmacist

Storage : Store below 30°C. Protect from moisture and light

Shelf-life : 3 years from date of manufacture

Presentation/Packing : Soft Capsules 400 I.U x 30's.
Soft Capsules 400 I.U x 50's, 90's, 500's (For export markets only)

Manufactured by : Hovid Bhd. 121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar), 30010 Ipoh, Perak.

Date of Revision : June 2023