

Important information. Please read carefully.

Asmafen™

Ketotifen 1mg

COMPOSITION

Each 5ml contains Ketotifen (as hydrogen fumarate) 1 mg. Parabens 0.15% w/v as preservative.

PHARMACODYNAMICS

Ketotifen is a sedating antihistamine that also has a stabilizing action on mast cells analogous to that of sodium cromoglicate.

PHARMACOKINETICS

Ketotifen fumarate is almost completely absorbed from gastrointestinal tract after oral doses, but bioavailability is reported to be only about 50% due to hepatic first-pass metabolism. Peak plasma concentrations occur 2 to 4 hours after an oral dose. It is mainly excreted in the urine as inactive metabolites with a small amount of unchanged drug. The terminal elimination half-life is about 21 hours.

INDICATIONS

Asmafen is indicated for long-term prevention of bronchial asthma, allergic bronchitis, asthmatic symptoms associated with hay fever. It is also indicated for the prevention and treatment of other allergic conditions such as allergic rhinitis, conjunctivitis and allergic skin disorders.

DOSAGE AND ADMINISTRATION

Unless otherwise prescribed by the physician.

Adults and children above 12 years	5 to 10ml (twice a day)
Children 2 to 12 years	5ml (twice a day)
Children 6 months to 2 years	2.5ml (twice a day)

CONTRAINDICATIONS

Asmafen is contraindicated for use during acute attacks of asthma and in patients who are on oral antidiabetic treatment

PRECAUTIONS

Do not discontinue treatment abruptly as this may lead to recurrence of the symptoms of asthma. Should withdrawal be necessary, reduce the dose gradually over a period of one week. Avoid alcoholic drinks. Use in pregnancy is best avoided.

When necessary, close medical supervision is required.

DRUG INTERACTIONS

Ketotifen may potentiate the sedative effects of other CNS depressants, antihistamines and alcohol.

SIDE EFFECTS

This preparation may cause drowsiness. Patients should be advised against driving motor vehicles or operating machinery. Other side effects such as dry mouth, dizziness and weight gain may also occur.

OVERDOSAGE AND TREATMENT

Overdosage may lead to drowsiness, confusion, dyspnoea, bradycardia or tachycardia, disorientation, and convulsions. If overdosage is recent, the stomach should be emptied by gastric lavage. Other supportive treatments should also be given if necessary.

PRODUCT DESCRIPTION, DOSAGE FORMS AND PACKAGING AVAILABLE

Clear, colourless syrup with peach flavour in packs of 120ml.

SHELF-LIFE

The expiry date is indicated on the packaging.

STORAGE

Protect from light. Store below 30°C.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date: 27-Dec-2013

Manufactured and Marketed by

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.

AS0403-001

