

SUNQUADIN LOZENGES 0.25MG

Each tablet contains:-

Dequalinium Chloride	0.25 mg
----------------------	---------

Product Description:

Round, flat, orange coloured and orange flavoured, chewable tablet with single score on one side and "SW" debossed.

Pharmacodynamics:

It has cationic germicidal surface active properties similar to other quaternary ammonium antiseptics. The mechanism of action of surface active agents is unknown, although numerous theories have been proposed. The anti-bacterial action appears to be related neither to the ability to denature protein nor to their ability to lower surface tension. Evidence has been presented that the major site of action of these compounds may be the cell membrane, where the agents may cause the escape of enzymes, coenzymes and metabolic intermediates.

Dequalinium is an anti-bacterial and anti-fungal agent, active against many gram-positive and gram-negative bacteria and against *Borrelia vincenti*, *Candida albicans* and some species of *Trichophyton*.

Pharmacokinetics:

No data available.

Indication:

It is indicated in the treatment of infections of the mouth and throat.

Recommended Dosage:

One tablet to be sucked slowly 2 to 3 hourly or as required.

Route of Administration:

Oral

Contraindications:

It is contraindicated in patients who have demonstrated hypersensitivity to it.

Warnings and Precautions:

Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea and vomiting may be serious. Therefore if sore throat persists for more than 2 days, discontinue use and consult physician. Allergic skin reactions have been known to occur.

Pregnancy and Lactation:

No data available.

Side Effects:

Due to low strength, side effects are rare. The possible side effects (if taken a large amount) are nausea and vomiting. Strong solution may cause oesophageal damage and necrosis. Allergic skin reactions may occur.

Interactions with Other Medicaments:

No known drug interaction.

Symptoms and Treatment of Overdose:

Due to its low strength, toxicity caused by overdosage is rare. Toxic symptoms include dyspnoea and cyanosis due to paralysis of the respiratory muscles, possibly leading to asphyxia. Depression of central nervous system (possibly preceded by excitement and with convulsions), hypotension and coma may also occur.

Milk, gelatin solution or egg white may be given; the use of mild soap solution has been suggested; respiration should be assisted and treatment for shock given. Convulsion can be controlled by intravenous injection of diazepam.

Effects on Ability to Drive and Use Machine: Not applicable.

Storage Conditions: Store at or below 30°C.

Pack size: Blister pack: A box of 10 x 10, 50 x 10 and 100 x 10 tablets per strip.

Pack size (export only): A bottle of 500 and 1000 tablets.

Shelf-life: 5 years.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer & Product Registration Holder:

SUNWARD PHARMACEUTICAL SDN. BHD.

No. 9, 11 & 17, Jalan Kempas 4,
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.

TL693A-R9
Revision Date: 25/08/2022