

VITAMIN B12 TABLET 50MCG

Each tablet contains:
Cyanocobalamin 50 mcg

Product Description:

Oval, yellow, sugar-coated tablets with no markings or embossment.

Pharmacodynamics:

The vitamin is essential for the normal functioning of all cells, but particularly for cells of the bone marrow, the nervous system and the gastro-intestinal tract. It appears to facilitate reduction reactions and participate in the transfer of methyl groups. Vitamin B12 is involved in protein, carbohydrate and fat metabolism, but its chief importance in mammalian tissues seems to be, together with folic acid, in the anabolism of deoxyribonucleic acid in all cells. Vitamin B12 acts in the form of its co-enzymes, in which the vitamin is linked to adenine and sugar, which catalyze specific reactions in intermediary metabolism. Vitamin B12 is a complex water soluble compound and is neutral. It is composed of two heterocyclic systems, a benzimidazole and a modified prophyryn nucleus.

Pharmacokinetics:

Cyanocobalamin is extensively bound to specific plasma proteins called transcobalamins; transcobalamin II appears to be involved in the rapid transport of cyanocobalamin to tissues. Cyanocobalamin is quickly taken up by the intestinal mucosa and held there for at least 2 to 3 hours. It does not appear in the blood or tissues for 2 to 3 hours and peak concentrations does not occur until 8 to 12 hours after the dose. It accumulates and reaches maximum concentration in the liver within 24 hours. By 5 to 6 days, 60 to 70% of the dose is in the liver. When given in the large therapeutic doses it may be irregularly absorbed and absorption is impaired in patients with an absence of intrinsic factor, with a malabsorption syndrome or with disease or abnormality of the gut or after gastrectomy.

Indication:

It is indicated in the treatment of cobalamin deficiency.

For treatment of pernicious anaemia when parenteral administration is not possible or not advised.

Recommended Dose:

In deficiency states, 50 to 150 mcg or as professionally prescribed.

In pernicious anaemia, 1000 mcg daily.

Route of Administration:

Oral

Contraindications:

It is contraindicated in patients who are sensitive to it.

Warnings and Precautions:

Cyanocobalamin should not be given before a diagnosis has been fully established because of the possibility of masking symptoms of subacute degeneration of the spinal cord. Cyanocobalamin is not a suitable form of vitamin B12 for the treatment of optic neuropathies associated with raised plasma concentrations of cyanocobalamin. The absorption of cyanocobalamin is reduced by metformin, potassium citrate and calcium

gluconate. Ascorbic acid destroys cyanocobalamin in-vitro. The destructive effect is affected by the concentration of both and the food source of cyanocobalamin.

Interactions with Other Medicaments:

Concomitant administration of antibiotics, methotrexate or pyrimethamine invalidate folic acid and vitamin B12 diagnostic blood assays.

Colchicine, para-aminosalicylic acid and heavy alcohol intake for longer than 2 weeks may produce malabsorption of vitamin B12.

Pregnancy and Lactation:

It is not known whether vitamin B12 can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Adequate and well-controlled studies have not been done in pregnant women. However, vitamin B12 is an essential vitamin and requirements are increased during pregnancy. Vitamin B12 appears in the milk of nursing mothers in concentrations which approximate the mother's vitamin B12 blood level.

Side Effects:

Cyanocobalamin is very free of side effects through occasional hypersensitivity reactions have occurred.

Symptoms and Treatment of Overdose:

No documented reports on the symptoms and treatment of overdosage available. Therapeutic or accidental overdosage resulting in adverse reactions or fatalities quite unlikely, considering its therapeutic dose and its being very well tolerated.

Effects on Ability to Drive and Use Machine:

Not known.

Storage Conditions: Store at or below 30°C. Protect from light.

Pack Sizes: Bottle pack: 180, 500 and 600 tablets.

Blister pack: 10x10, 50x10 and 100x10 tablets.

Shelf Life: 2 years.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer & Product Registration Holder:

SUNWARD PHARMACEUTICAL SDN. BHD.

No. 9,11 & 17, Jalan Kempas 4,

Taman Perindustrian Tampoi Indah,

81200 Johor Bahru, Johor, Malaysia.

TL704C-R5
Revision Date: 01/09/2022