

[싱가포르][말레이시아]
폴리바마이신연고 설명서
Size : 105 × 200 (mm)

- 7/1 수정사항
- 1. Zypas 주소 추가
 - 2. Date of Revision 추가
 - 3. 공간확보 위한 전체 Layout 수정

- 8/14 수정사항
- 1. Zypas 사명 변경

Antibiotic Compound
POLYBAMYCIN ointment
Bacitracin, Neomycin, Polymyxin B Oint

【DESCRIPTION】
Each gram contains : Bacitracin 400 units, neomycin sulfate 3.5mg(potency), polymyxin B sulfate 5,000 units ; special white petrolatum q.s.

【ACTIONS】
The overlapping spectra of the antibiotics provide effective antibacterial action against most commonly occurring bacteria known to be topical invaders. The range of antibacterial activity encompasses many bacteria which are, or have become, resistant to other antibiotics or chemically—notably Pseudomonas and Staphylococcus. In susceptible organisms, resistance rarely develops, even on repeated or prolonged use.

【INDICATIONS】
Infected burn due to susceptible organisms, primary pyoderma (impetigo, ecthyma, sycosis vulgaris, paronychia), secondarily infected dermatitis(eczema, herpes and seborrheic dermatitis, infection after surgical incisions and skin grafts), traumatic lesions, purulence, external otitis.

【DOSAGE AND ADMINISTRATION】
A small amount should be applied to the affected areas once to several times daily.

【ADVERSE EFFECTS】
Skin allergic disorders, ototoxicity, and deafness in patients with extensive skin damage or perforated tympanic membranes.

【CONTRAINDICATIONS/PRECAUTIONS】
1. Avoid long term use as it may lead to skin sensitization.
2. It should not be given to patients with a history of hypersensitivity to any of the components of this ointment.

【USAGE IN PREGNANCY】 No known side effects

【INCOMPATIBILITIES/DRUG INTERACTIONS】
Neomycin sulfate : It is incompatible with beta-lactam antibiotics such as carbenicillin sodium in vivo. These antibiotics should be administered separately when both are required.
Polymyxin B sulfate : Loss of activity has been reported with many antimicrobial agents including amphotericin, ampicillin sodium, cephalothin sodium, cephalozin sodium, chloramphenicol sodium succinate, nitrofurantoin sodium and tetracycline hydrochloride ; and also with calcium and magnesium salts, heparin sodium, and prednisolone sodium phosphate. It is rapidly inactivated by strong acids and alkalis.

【PACKING SIZE】 10 g/Tube

【STORAGE】 Store in a tight container at temperature not exceeding 30℃.

【EXPIRY】 2 years from manufacturing date.

Product Registration Holder :
The Zypas Medical Co.
No. 15, Jalan Mega 1/5,
Taman Perindustrian Nusa Cemerlang, 79200 Iskandar Puteri Johor, Malaysia

Date of Revision : 2024.07.01

