

AMENDED PACKAGE INSERT

TRICONAZOLE CREAM

Composition:

Triamcinolone Acetonide BP	0.1% w/w
Econazole Nitrate BP	1.0% w/w

Pharmacology:

Triamcinolone acetonide and econazole nitrate combination has anti-inflammatory, anti-pruritic, anti-fungal and anti-bacterial action.

Econazole acts by damaging cell membranes. The permeability of the fungal cell is increased. Subcellular membranes in the cytoplasm are damaged. The site action is most probably the unsaturated fatty acid acyl moiety of membrane phospholipids.

Triamcinolone acetonide is a more potent form of the naturally occurring adrenocortical hormone. The mechanism of anti-inflammatory activity of the corticosteroids is unclear, but there is some evidence to suggest that a recognizable correlation exists between vasoconstrictor potency and therapeutic efficacy in man.

When applied topically, particularly to large areas, the triamcinolone acetonide may be absorbed in sufficient amount to cause systemic effect. It is slowly metabolized in the skin and thus tend to be systemically absorbed to a greater extent.

After topical application to the skin of normal subjects, systemic absorption of econazole nitrate is extremely low. Although most of the applied drug remains on the skin surface, drug concentrations were found in the stratum corneum which, by far, exceeded the minimum inhibitory concentration for dermatophytes. Inhibitory concentrations are achieved in the epidermis and as deep as the middle region of the dermis. Less than 1% of the applied dose was recovered in the urine and faeces.

Indications:

Inflammatory dermatoses with mycotic or bacterial infection. Inflammatory dermatomycoses caused by dermatophytes, yeasts and molds.

Contraindications:

It is contraindicated in patient who are hypersensitive to econazole nitrate, triamcinolone or other corticosteroids.

Dosage:

Route of Administration : Topical

Apply cream to the affected areas twice daily.

Warning and Precautions:

The medication should be discontinued if irritation occurred.

Do not use in and around eyes.

Special precaution should be taken when medication is used in children.

Use in Pregnancy and Lactation:

Pregnant women and nursing mothers should seek the advice of the physician before using this product.

Interactions with Other Medications

Econazole is a known inhibitor of CYP3A4/2C9. However, due to the limited systemic availability after cutaneous application, clinically relevant interactions are unlikely to occur, but have been reported for oral anticoagulants. In patients taking oral anticoagulants, such as warfarin or acenocoumarol, caution should be exercised and the anticoagulant effect should be monitored.

Side Effects/Adverse Effect:

The side effects consist of itching, burning, stinging, erythema, dryness, folliculitis, hypertrichosis, acne form eruption, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, miliaria.

Symptoms and Treatment for Overdosage and Antidote(s):

Topically applied corticosteroids can be absorbed in sufficient amount to produce systemic effect.

(See Precaution/Warning).

Pack size: A tube of 15g.

A box of 50 x 15g / tube

Pack size (export only): A jar of 400g.

Shelf-life: 3 years.

Storage conditions: Store at or below 30°C. Protect from light.

Description: A white and smooth cream.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer & Product Holder:

SUNWARD PHARMACEUTICAL SDN. BHD.

No. 9, 11 & 17, Jalan Kempas 4,

Taman Perindustrian Tampoi Indah,

81200 Johor Bahru, Johor, Malaysia.

EL225A-R8

Revised Date 11/12/2020