



212/15243278/0817

Directions for use

Read carefully!

Composition

Each 100 ml contains:	
Sodium Chloride	0.23 g
Glucose Monohydrate for Parenteral Use	4.13 g
Water for Injections to	100 ml
Electrolytes:	mmol/l (mEq/l)
Na ⁺	40
Cl ⁻	40
Caloric value:	640 kJ/l $\triangle$ 150 kcal/l
Osmolarity:	288 mOsm/l

Characteristics

B. Braun 0.23% Sodium Chloride And 3.75% Glucose Intravenous Infusion B.P. represents an isotonic i.v. solution.

The outstanding feature of this solution is the markedly low electrolyte content which makes it particularly suitable for use in case of hypertonic dehydration. Since the young are known to require less electrolytes than adults, the solution is ideally suited for use in pediatrics.

Mechanisms of action

Glucose is absorbed from the gastrointestinal tract, and oxidised as a source of energy, or stored in the liver as glycogen. It is the only energy substrate which is directly, instantly and universally utilized by the body.

Glucose is vital for the myocardium, brain and nerves. Sodium Chloride is the principal salt involved in maintaining the osmotic tension of the blood and tissue.

Changes in sodium and chloride levels change this osmotic tension and hence influence the movement of fluids and diffusion of salts in cellular tissue.

Indications

Hypertonic (esp. hypernatremic) dehydration,
For electrolyte maintenance in pediatrics,
Caloric supply,
Vehicle solution for supplementary medication.

B. Braun 0.23% Sodium Chloride And 3.75% Glucose Intravenous Infusion B.P.

Dosage

According to individual requirements approximately 1000 ml/day
Drop rate : 120-180 drops/min corresponding to 360-540 ml/h

Route of administration I.V.

Contraindications

Hyperhydration, oedema
Hypotonic dehydration
Cardiac insufficiency
Diabetes mellitus

Precautions

Use with caution in patients with hypertension and diabetes mellitus. Blood glucose, serum electrolytes and water balance should be monitored regularly.

The compatibility of any additives to this solution should be checked before use.

Side effects

Hyperglycaemia and glycosuria may occur if glucose tolerance is impaired. These can be avoided by reducing the dosage and infusion rate or by administering insulin.

Symptoms and treatment for overdose

If hyperglycaemia or glycosuria occurs, the dosage and infusion rate should be reduced, or insulin may be administered.

Shelf life

The product must not be used beyond the expiry date stated on the label.

Storage

The product should not be stored above the temperature stated on the label.

Presentation

500 ml plastic container.

Product registration holder and
manufactured by:

B. BRAUN B. Braun Medical Industries Sdn. Bhd.
11900 Bayan Lepas, Penang, Malaysia.

Approval for Printing

B. BRAUN Melsungen AG

Approved for Printing ☐

Approved for Printing
when corrected ☐

New draft required ☐

Date _____ Signature _____

Name in capital letters

schwarz

Format = 148 x 210 mm (DIN A5)
1 Seite

Lätus



4311

MY_212

212/15243278/0817
GIF (EP)
Standort Penang

Font size: 9 pt.

G 170375

