


CIMULCER TABLETS

Cimetidine Tablets



COMPOSITION

CIMULCER TABLETS 200 mg

Each tablet contains: Cimetidine 200 mg

CIMULCER TABLETS 400 mg

Each tablet contains: Cimetidine 400 mg

DOSAGE FORM AVAILABLE

Tablet

PRODUCT DESCRIPTION

CIMULCER TABLETS 200 mg

Round, biconvex, green film-coated tablet, impressed BIOLAB's logo on one face

CIMULCER TABLETS 400 mg

Oblong, biconvex, green film-coated tablet, impressed "BIOLAB" on one face and "400" on the other face

ACTIONS

Pharmacological actions:

CIMULCER (Cimetidine) is a histamine H₂-receptor antagonist.

Mechanism of action:

Cimetidine competitively inhibits the action of histamine on the H₂-receptors of parietal cells, reducing gastric output and concentration under basal conditions and also stimulated by food, insulin, histamine, pentagastrin and caffeine.

Pharmacokinetics:

Absorption: Cimetidine is rapidly and well absorbed following oral administration.

A small portion of orally administered is metabolized during first pass through the liver, resulting in an average bioavailability of 60-70% when compared to IV route.

Distribution: Cimetidine is widely distributed throughout the body and is 15-20% bound to plasma proteins. It is distributed into milk.

Elimination: Cimetidine is metabolized in the liver to sulfoxide and 5-hydroxy methyl derivatives.

It has a plasma half life about 2 hours in patients with normal renal function, 2.9 hours in patients with creatinine clearance of 20-50 ml per minute, 3.7 hours in patients with creatinine clearances of less than 20 ml per minute and 5 hours in anephric patients. It is excreted in urine within 24 hours.

INDICATIONS

Cimulcer is a histamine H-2 receptor antagonist. Its main action is to inhibit gastric acid secretion. It is used in gastric and duodenal ulcer, and in other conditions associated with gastric hyperacidity i.e. Zollinger-Ellison Syndrome.

DOSAGE AND ADMINISTRATIONS

Gastric and Duodenal Ulcer:

Adults: Oral, 200 mg 3 times daily after meals and 400 mg at bedtime or 400 mg twice daily at breakfast and at bedtime or 800 mg once daily at bedtime for 4-6 weeks or as directed by the physician.

Zollinger-Ellison Syndrome:

Adults: Oral, 200-400 mg, 3-4 times a day after meals and 400 mg at bedtime.

OVERDOSAGE

A patient who took about 12 g of Cimetidine had slurred speech, high pulse-rate, and dilated pupils. Five hours after gastric lavage he was confused, agitated, disorientated and making staccato nonsensical conversation. He recovered from these mental symptoms the next day. There were no changes in ECG, blood indexes, liver or kidney.

CONTRAINDICATIONS

Cimetidine is contraindicated in the patients who are known to have hypersensitivity to the drug.

PRECAUTIONS AND WARNINGS

CIMULCER should not be used in pregnancy and lactation. It should be given with care and in reduced dosage to patients with impaired renal function.

PREGNANCY AND LACTATION

There has been limited experience to date using cimetidine in pregnant women and the drug should be used in pregnant patients only when clearly needed.

Since cimetidine is distributed into milk and as a general rule, nursing should not be taken during therapy with the drug.

SIDE EFFECTS

Mild diarrhoea, muscle pain, dizziness, and skin rash may occasionally occur.

DRUG INTERACTIONS

Cimetidine, as enzyme inhibitor of hepatic microsomal enzyme system, reduces the hepatic metabolism of some drugs including coumarin anticoagulant, phenytoin, propranolol, some benzodiazepines, lidocaine, theophylline, metronidazole, and triamterene, thereby decreasing elimination and increasing plasma concentration of these drugs bioavailability of drugs with high hepatic extraction ratios.

STORAGE

Store at temperature not exceeding 30°C. Protect from light.

Keep medicine out of reach of children.

Jauhkan daripada kanak-kanak.

The shelf-life period is 5 years.

SUPPLY

CIMULCER TABLETS 200 mg

Blister pack 50x10's

CIMULCER TABLETS 400 mg

Blister pack 10x10's

Date of revision
Sept 2017



BIOLAB

NYI(MMSB): R4(NOD_V/C 2-3/2014)
RAP(E)_TU

Manufactured by:

BIOLAB CO., LTD.

625 Bangpoo Industrial Estate,

Samutprakarn, Thailand

EET 101/A

