



File Name: Leaflet Cough Relief [Spore]

Received : 02.10.2018  
Amended : 02.04.2021

Output : -  
Request : Shuying  
Designer : Syafiq

Opened : 223w x 141h MM  
Closed : -  
Finished: -  
Colour : 1C

K



## THREE LEGS COUGH RELIEF UBAT BATUK CAP KAKI TIGA 三脚标止咳水

### THREE LEGS COUGH RELIEF

**Composition:** Each teaspoon (5ml) contains 100mg Guaifenesin, 5mg Sodium Benzoate

**Product Description:** Dark brown coloured, moderately viscous solution with a sweet, cooling peppermint flavour.

**Indication:** As an expectorant for the symptomatic relief of cough due to colds.

#### Dosage:

Adults: 10ml - 20 ml or 2 - 4 teaspoons  
6-12 years: 5ml - 10ml or 1 - 2 teaspoons  
2-6 years: 2.5ml - 5ml or ½ - 1 teaspoon  
To be taken every 4 hours or use as directed by your doctor.

**Route Of Administration:** Oral

**Contraindications:** Hypersensitivity to Guaifenesin

**Warnings And Precautions:** May cause drowsiness in some case. Do not administer to children below 2 years without supervision of doctor. Refrain from activities requiring alertness, such as driving vehicle or operating machine if affected.

If symptoms persist, please consult doctor.

Keep out of reach of children.

#### Interactions With Other Medicaments:

##### Diagnostic Interference:

Guaifenesin has been shown to produce a colour interference with certain clinical laboratory determinations of 5-hydroxyindoleacetic acid (5-HIAA) and vanilmandelic acid (VMA).

Guaifenesin could interfere with diagnosis of carcinoid syndrome. Patients being evaluated for carcinoid syndrome should therefore discontinue any preparation containing guaifenesin for 24 hours before collection of urine specimens for the determination of 5-HIAA.

#### Statement On Usage During Pregnancy And Lactation:

Available evidence and/or expert consensus is inconclusive or is inadequate for determining infant risk when used during pregnancy and breastfeeding. Weigh the potential benefits of drug treatment against potential risks before taking this drug during pregnancy and breastfeeding.

#### Adverse Effects/ Undesirable Effects:

Gastrointestinal discomfort, nausea and vomiting have occasionally been reported with guaifenesin, particularly in very large doses. Others include dizziness, diarrhea, headache and rash (including urticarial).

#### Symptoms And Treatment Of Overdose:

Nausea and vomiting may occur.

In the event of overdose, discontinue medication and seek medical help immediately.

#### Pharmacodynamics:

Guaifenesin is an expectorant which is thought to act by irritating the gastric mucosa and subsequently stimulating respiratory tract secretions. This increase in fluid increase the volume and decrease the viscosity of bronchial secretions.

#### Pharmacokinetics:

Guaifenesin is well absorbed from the gastrointestinal tract. It is metabolized and then excreted in the urine. The half- life in plasma is approximately 1 hour.

**Effects On Ability To Drive And Use Machines:** Not known

**Storage Conditions:** Store below 30°C. Protect from light and moisture.

**Shelf Life:** 5 years

**Dosage Form:** Liquid

**Packing Size:** 60ml, 80ml, 100ml and 120ml

The information contained in this leaflet is limited. For further information, please consult your pharmacist or doctor.

### UBAT BATUK CAP KAKI TIGA

**Komposisi:** Setiap sudu (5ml) mengandungi 100mg Guaifenesin, 5mg Natrium Benzoat

**Deskripsi:** Larutan berwarna coklat pekat, kurang likat dengan rasa manis dan pudina segar.

**Indikasi:** Sebagai ekspektoran untuk melegakan simptom batuk akibat selsema.

#### Sukatan:

Dewasa: 10ml - 20 ml atau 2 - 4 sudu teh  
6-12 tahun: 5ml - 10ml atau 1 - 2 sudu teh  
2-6 tahun: 2.5ml - 5ml atau ½ - 1 sudu teh  
Ulangi sukatan setiap 4 jam atau seperti yang disarankan oleh doktor anda.

**Cara Pengambilan Ubat:** Oral

**Kontraindikasi:** Hipersensitiviti terhadap Guaifenesin.

#### Amaran Dan Perhatian:

Boleh menyebabkan rasa mengantuk dalam sesetengah keadaan. Jangan berikan kepada kanak-kanak di bawah umur 2 tahun tanpa pengawasan doktor. Elakkan diri daripada aktiviti yang memerlukan tumpuan, seperti memandu kenderaan atau mengendalikan mesin.

Jika simptom berlarutan, sila rujuk kepada doktor.

Jauhkan daripada capaian kanak-kanak.

**Interaksi Dengan Ubat Lain:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Kehamilan Dan Penyusuan:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Kesan Sampingan:** Ketidakselesaan gastrousus, loya dan muntah juga dikaitkan dengan guaifenesin, terutamanya dalam dos yang besar. Termasuk pening, cirit-birit, sakit kepala dan ruam (termasuk urtikaria).

#### Symptom Dan Rawatan Untuk Dos Berlebihan:

Loya dan muntah boleh berlaku.

Sekiranya melebihi dos, hentikan ubat dan dapatkan bantuan perubatan dengan segera.

**Farmakodinamik:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Farmakokinetik:** Sila rujuk kepada penerangan Bahasa Inggeris.

**Kesan Ke Atas Keupayaan Memandu Dan Penggunaan Mesin:** Tiada pengesahan

**Keadaan Penyimpanan:** Simpan di bawah 30°C. Lindungi dari cahaya dan kelembapan.

**Tempoh Hayat:** 5 tahun

**Bentuk Dos:** Cecair

**Saiz Bungkusan:** 60ml, 80ml, 100ml dan 120ml

Maklumat yang terkandung di dalam risalah ini adalah terhad. Untuk maklumat yang terperinci, sila hubungi ahli farmasi atau doktor.

### 三脚标止咳水

**成份:** 每茶匙(5毫升)含有100毫克愈创甘油醚和5毫克苯甲酸钠

**外观:** 深褐色液体, 带有粘性, 甜味和冰凉薄荷味。

**主治:** 作为缓解感冒咳嗽症状的祛痰剂。

#### 服量:

成人: 10毫升 - 20毫升或 2 - 4 茶匙

6-12 岁: 5毫升 - 10毫升或 1 - 2茶匙

2-6岁: 2.5毫升 - 5毫升或 ½ - 1茶匙

每隔四小时服用一次或遵照医生的指示服用。

**给药途径:** 口服

**禁忌:** 对愈创甘油醚过敏者。

**警告及注意:** 在某些情况下可能会引起嗜睡。若没有医生监督, 不要给予2岁以下的儿童使用。避免需要警觉性的活动, 如驾驶车辆或操作机器。

如症状持续, 请咨询医生。

远离儿童。

**药物相互作用:** 请参阅英文版本。

**怀孕和哺乳:** 请参阅英文版本。

**副作用:** 使用愈创甘油醚时可能会引起胃肠道不适, 恶心和呕吐, 特别是在非常大的剂量下。其他包括头晕, 腹泻, 头痛和皮疹 (包括荨麻疹)。

#### 药物过量症状及治疗:

可能会出现恶心和呕吐。

如果服用过量, 请立即停止服药并寻求医疗帮助。

**药效学:** 请参阅英文版本。

**药代动力学:** 请参阅英文版本。

**对驱动和使用机器的能力的影响:** 未知

**贮存:** 储藏低于30°C。避免阳光直射及潮湿处。远离儿童。

**有效期:** 5年

**剂型:** 液体

**包装:** 60毫升, 80毫升, 100毫升和120毫升

此说明书仅属概述, 欲知更详细资料, 可咨询您的药剂师或医生。

Manufactured by Product Registration Holder

**Syarikat Wen Ken Drug Sdn.Bhd.**

24, Jalan Lambak, Taman Johor,  
81200 Johor Bahru, Johor, Malaysia.

Singapore Distributor/ Pengedar Singapore/ 新加坡代理商:

**Wen Ken Marketing (S) Pte. Ltd.**

2, Alexandra Road, #02-08 Delta House,  
Singapore 159919.

Date of Revision: 02.04.2021