



RASITIN TABLET 10mg

Description: White, oval, top-scored tablet.

Content

Cloperastine HCl 10mg/tablet

Pharmacology

Cloperastine is an antitussive agent with an effect similar to that of codeine, it also has some antihistaminic effect.

Indications

For the relief of unproductive coughs

Adverse Effects

Dry mouth and drowsiness may occur.

Precautions

- Caution when driving or doing jobs requiring alertness because of possible drowsiness or dizziness.
- Safety for use in pregnancy and lactation has not been established.

Dosage and Administration

Adults - Oral, 10 to 20mg (1 to 2 tablets) three times a day.

Children 4 to 6 years - Oral, 2.5 - 5.0mg three times a day.

7 to 15 years - Oral, 5 to 10mg (1/2 to 1 tablet) three times a day Or as directed by physician.

Storage

: Keep container tightly closed.

Store in a cool, dry place, below 30C. Protect from light.

Presentation/Packing : Tablet 10mg x 500's

Shelflife : 3 years

Manufactured & : **PRIME PHARMACEUTICAL SDN. BHD.**

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

27/02/2018