

HOVID-CIMETIDINE FILM-COATED TABLET 400 MG

VICIM07-var (MY)

DESCRIPTION

Oblong, pale yellow film-coated tablet, convex faces and break-bar embossed on one face.

COMPOSITION

Each tablet contains: Cimetidine 400 mg.

PHARMACODYNAMICS

Cimetidine inhibits basal and nocturnal gastric acid secretion by competitive inhibition of the action of histamine at the histamine H₂-receptors. It also inhibits gastric acid secretion stimulated by food, betasole, pentagastrin, caffeine, insulin and physiological vagal reflex. It is selective in its actions and is virtually without influence on those effects of histamine that are mediated through H₂-receptors.

PHARMACOKINETICS**Absorption**

Cimetidine is well absorbed (about 60%) by the oral route.

Half-life

Plasma half-life of cimetidine is approximately 2 hours.

Protein binding

Cimetidine is weakly bound, about 20% to plasma proteins.

Distribution

Cimetidine crosses the placental barrier and is excreted in breast milk. It is distributed widely in almost all tissues except the brain.

Metabolism

Cimetidine is partially metabolized in the liver.

Excretion

Most of an oral dose of cimetidine is excreted unchanged in the urine within 24 hours. Some is excreted in the bile.

INDICATIONS

Cimetidine is indicated in the:

- prophylaxis and treatment of duodenal ulcer.
- short-term treatment of active benign gastric ulcer.
- treatment of pathological gastric hypersecretion states, such as Zollinger-Ellison syndrome.
- treatment of acute gastroesophageal reflux disease.
- treatment of upper gastrointestinal bleeding.

CONTRAINDICATIONS

- Avoid in patients known to be hypersensitive to histamine H₂-receptors antagonists.
- It should not be used in nursing mothers.

PRECAUTIONS

- Caution in patients with cirrhosis, history of portal systemic encephalopathy, hepatic and renal function impairment.
- Safety for use in pregnancy has not been established.
- Dosage should be reduced in patients with impaired renal function.
- Care should be taken in patients with a history of peptic ulcer, particularly the elderly.

PREGNANCY AND LACTATION

Safety for use in pregnancy and lactation has not been established.

MAIN SIDE/ADVERSE EFFECTS

Diarrhoea, dizziness, tiredness, rashes. Reversible confusional states, especially in the elderly or in seriously ill patients such as those with renal failure, gynaecomastia and impotence have occasionally occurred.

DRUG INTERACTIONS

Cimetidine can prolong the elimination of drugs metabolised by oxidation in the liver.

Although pharmacological interactions between cimetidine and a number of drugs have been demonstrated e.g. diazepam and propranolol, only those with oral anticoagulants, phenytoin, theophylline and intravenous lidocaine appear, to date, to be of clinical significance. Close monitoring of patients on cimetidine receiving oral anticoagulants or phenytoin is recommended and a reduction in the dosage of these drugs may be necessary.

In patients on drug treatment or with illnesses that could cause falls in blood cell count, the possibility that H₂-receptor antagonism could potentiate this effect should be borne in mind.

Cimetidine has the potential to affect the absorption, metabolism or renal excretion of other drugs which is particularly important when drugs with a narrow therapeutic index are administered concurrently. The altered pharmacokinetics may necessitate dosage adjustment of the affected drug or discontinuation of treatment.

Interactions may occur by several mechanisms including:

- 1) Inhibition of certain cytochrome P450 enzymes (including CYP1A2, CYP2C9, CYP2D6 and CYP3A3/A4, and CYP2C18); Inhibition of these enzymes may result in increased plasma levels of certain drugs including warfarin-type coumarin anticoagulants (e.g. warfarin), tricyclic antidepressants (e.g. amitriptyline), class I antiarrhythmics (e.g. lidocaine), calcium channel blockers (e.g. nifedipine, diltiazem), oral sulfonylureas (e.g. glipizide), phenytoin, theophylline and metoprolol.
- 2) Competition for renal tubular secretion; This may result in increased plasma levels of certain drugs including procainamide, metformin, ciclosporin and tacrolimus.
- 3) Alteration of gastric pH; The bioavailability of certain drugs may be affected. This can result in either an increase in absorption (e.g. atazanavir) or a decrease in absorption (e.g. some azole antifungals such as ketoconazole, itraconazole or posaconazole).
- 4) Unknown mechanisms; Cimetidine may potentiate the myelosuppressive effects (e.g. neutropenia, agranulocytosis) of chemotherapeutic agents such as carmustine, fluorouracil, epirubicin, or therapies such as radiation. Isolated cases of clinically relevant interactions have been documented with narcotic analgesics (e.g. morphine).

OVERDOSAGE**Clinical features:**

Allergic reaction, bradycardia or tachycardia, bronchospasm, confusion, neutropenia or other blood dyscrasias and fever may occur.

Treat overdosage by emesis or gastric lavage; artificial respiration in the event of respiratory failure and administration of beta-blocker to control tachycardia.

DOSAGE AND ADMINISTRATION

Adults (Oral).

Duodenal or benign gastric ulcer:

A single daily dose of 800 mg at bedtime, or 400 mg twice a day with breakfast and at bedtime, or 200 mg three times a day with meals and 400 mg at bedtime (1.0 g/day), or 400 mg four times a day with meals and at bedtime (1.6 g/day).

Treatment should be given initially for at least four weeks (six weeks in benign gastric ulcer).

Oesophageal reflux disease:

400 mg four times a day, with meals and at bedtime, for 4 to 8 weeks.

Gastric hypersecretion states (eg Zollinger-Ellison Syndrome):

400 mg four times a day.

The total daily dose should not exceed 2.4 g. Dosage should be reduced in patients with impaired renal function. Antacids can be made available to all patients until symptoms disappear.

Children (more than one year old):

Oral, 25 - 30 mg/kg of body weight per day in divided doses.

Storage:

Store below 30°C. Protect from light and moisture.

Presentation/Packing:

Blisters of 10 x 10's
100's, 500's, 1000's (For export only)
(Not all presentations are available locally)

Product Registration Holder: HOVID Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Manufactured by: HOVID Bhd.

Lot 56442, 7 1/2 Miles, Jalan Ipoh / Chemor,
31200 Chemor, Perak, Malaysia.

Revision date: May 2021