

SELFIDE LOTION

DESCRIPTION

A reddish-brown suspension containing 2.5g of selenium sulfide per 100ml.

INDICATION

For the treatment of seborrheic dermatitis, tinea versicolor, dandruff and as an adjunct in the treatment of tinea capitis.

RECOMMENDED DOSE

SCALP TREATMENT - Apply Selfide lotion to wet hair, lather and leave on scalp for 3 minutes. Rinse. Repeat treatment. Then rinse thoroughly in running water. Use twice weekly at first, then as necessary as physician directs.

SKIN TREATMENT - Apply undiluted to affected areas for 10 minutes. Rinse thoroughly. Repeat daily for 1 week or as physician directs.

ROUTE OF ADMINISTRATION

For topical use.

CONTRAINDICATIONS

- Hypersensitivity to any of the ingredients
- Do not allow contact with broken and severely inflamed skin

WARNINGS AND PRECAUTIONS

- For external use only. Keep out of eyes; if Selfide lotion gets into the eyes, wash thoroughly with cold water.
- Do not leave the lotion in contact with the hair or skin for more than the recommended duration as irritation, burning sensation or blistering may occur and do not use more often than recommended.
- Selfide lotion is not to be ingested.
- In the event of accidental ingestion, take all appropriate measures immediately.
- Selfide lotion must not be applied to or on infected or broken skin as this may cause systemic absorption of the selenium sulfide.

PREGNANCY AND LACTATION

Avoid during pregnancy and breastfeeding.

SIDE EFFECTS

Immune system disorder

Frequency unknown: Hypersensitivity, rash and urticarial

Skin and subcutaneous tissue disorder

Frequency unknown: Irritation and sensitization, sometimes described as a burning sensation, blistering can occur, especially if the shampoo is kept in contact with hair or skin for longer than the recommended duration.

Alopecia can occur. Hair colour changes may occur; this can be avoided or minimized by thorough washing of the hair after treatment.

Seborrhea (oiliness of hair and scalp) or application site dryness may occur.

INTERACTION WITH OTHER MEDICAMENTS

- If the hair has been damaged by chemical substances (like dyeing or bleaching products or as a result of a permanent wave or relaxing treatment) or if the hair is grey or white, discoloration of the hair may occur. In order to avoid discoloration of the hair it must be thoroughly rinsed immediately after use.
- Selfide lotion should be very thoroughly rinsed from hair before dyeing, tinting, or permanent waving the hair. It should not be applied for a period of two days before or after any of these procedures.
- Gold, silver and other metallic jewellery should be removed prior to use of Selfide lotion, since discoloration of the metals may occur.

SYMPTOMS AND TREATMENT OF OVERDOSE

If ingested, vomiting should be provoked and general supportive measures given.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

No studies on the effect on the ability to drive or use machines have been performed.

PHARMACODYNAMICS

Selenium sulphide appears to have a cytostatic effect on cells of the epidermis and follicular epithelium, thus reducing corneocyte production. Selenium sulphide acts as an antiseborrheic agent which effectively controls itching and scaling dandruff. It has activity against certain dermatophytes including *Pityrosporum orbiculare*, the organism causing pityriasis versicolor (tinea versicolor).

EXCIPIENT

Titanium dioxide, Bentonite, Citric acid, Sodium dihydrogen phosphate, Sodium chloride, Colour perfume, Texapon 103L, Comperan LD, Cutina AGS, Dehyton K, Cetiol HE, Brunidox L.

PACKING AND STORAGE CONDITION

Bottle of 60 or 120ml.

Store in a cool dry place, below 30°C.

Keep out of reach of children.

IF SWALLOWED SEEK MEDICAL ADVICE IMMEDIATELY.

Shelf life: 3 years

MANUFACTURED BY:

T.O.Chemicals (1979) Ltd.

280, Soi Sabaijai, Suthisarnwinijai Road,

Samsennok, Huay-Kwang,

Bangkok 10310, Thailand

Revision date: 18 June 2020