

selsun yellow_12cm x 8cm

Dosage form and packaging available 30ml, 60ml, 120ml HDPE bottle Manufactured by INTERTHAI PHARMACEUTICAL MANUFACTURING LTD Bangkok, Thailand. Product Registration Holder Rohto-Mentholatum (M) Sdn Bhd (354691-U), Unit 9-1, Level 9, Wisma Averis, Avenue 5, Bangsar South City, No.8 Jalan Kerinchi, 59200 Kuala Lumpur. This leaflet was last revised in September 2020.	 MENTHOLATUM® selsun® 2.5 % SUSPENSION Selenium Sulfide 2.5% Selsun is a registered trademark
---	---

Read all of this leaflet carefully before you start using this medicine • Keep this leaflet. You may need to refer to it again. • Should there be any further questions, please refer to your doctor/pharmacist. • This medicine has been prescribed for you only. Do not pass it on for usage to others. It may be harmful to them, even if their signs of illness are the same as yours. • In the event of any side effects developed after using the medicine, talk to your doctor or pharmacist. In this leaflet 1. What Selsun 2.5% Suspension is and what it is used for 2. What you need to know before you use Selsun 2.5% Suspension 3. How to use Selsun 2.5% Suspension 4. How to store Selsun 2.5% Suspension 5. Contents of the pack and other information 1. What Selsun 2.5% Suspension is and what it is used for Pharmacodynamic Selenium sulphide appears to have a cytostatic effect on cells of the epidermis and follicular epithelium, thus reducing corneocyte production. Selsun acts as an antiseborrhoeic agent which effectively controls itching and scaling dandruff. It has activity against certain dermatophytes including Pityrosporum obliculare, the organism causing pityriasis versicolor (tinea versicolor).	2. What you need to know before you use Selsun 2.5% Suspension Indication For treatment of Seborrhoeic Dermatitis, Tinea Versicolor, Dandruff and as an adjunct in the treatment of Tinea Capitis. Contraindication Not to be used by patients allergic to ingredients of this product. Caution • For external use only, do not swallow and avoid contact with eyes (Elakkan daripada terkena mata). Keep out of reach of children. (Jauhi daripada pencapalan kanak-kanak) • Do not use on broken or inflamed skin. (Selenium sulphide tidak boleh digunakan pada kulit yang pecah dan radang). Skin irritation or discoloration of hair may occur and can be minimized or avoided by rinsing thoroughly. • If skin irritation persists, discontinue use. • Not recommended for use by patients allergic to the ingredients, on infants and for treatment of tinea versicolor in pregnant women. • Avoid use 48 hours before or after permanent waving, tinting or bleaching. Avoid contact with jewellery. Pregnancy and Lactation There are no adequate data from the use of Selenium Sulfide in pregnant women. Animal studies are insufficient with respect to effects on pregnancy, foetal development and postnatal development. Avoid during pregnancy and breastfeeding.	Symptoms of Overdose and Treatment If swallowed, vomiting should be provoked or contact a doctor or poisons information centre. 3. How to use Selsun 2.5% Suspension Route of Administration Topical Recommended Dosage A. Scalp Treatment 1. Apply Selsun to wet hair, lather and leave on scalp for 3 minutes. 2. Rinse 3. Repeat treatment then rinse thoroughly in running water. 4. Use twice weekly at first, then as necessary or as doctor directs. B. Skin Treatment 1. Apply undiluted to affected areas for 10 minutes. 2. Rinse thoroughly 3. Repeat daily for 1 week or as doctor directs. 4. How to store Selsun 2.5% Suspension Storage Condition • Store below 30°C. 5. Contents of the pack and other information Product Description Beige (pale yellow) smooth uniform suspension with faint pleasant aroma, free from sulfide odour. Contents/ Mengandung DMDM Hydantoin 0.5% w/v Active Ingredient SELENIUM SULFIDE 25mg/ml Controlled medicine / Ubat Terkawal
--	---	--