
UNI-PAIN OINTMENT

Name and Strength of Active Substance:

Each 100 gm contains:
Methyl Salicylate B.P. 25 gm

Product Description:

White to pale yellow colour ointment with a characteristic odour.

Indication:

For temporary relief of minor muscular and rheumatic pains.

Application:

Apply a little to the affected part and gently massage. To be applied 3 or 4 times a day.

Contraindications:

Allergic reaction to any of the components of the preparation.
This product is contraindicated in children below 2 years of age.
Caution must be exercised when older children are treated.

Warning and precautions:

- i) For external use only.
- ii) If condition worsen, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.
- iii) Do not apply to wound or damaged skin.
- iv) Do not apply over large areas of the body or bandage.
- v) This product contains methyl salicylate and when applied or rub to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application of the skin for muscle or joint pain may increase the chances of bleeding.
- vi) It is dangerous to place any camphor containing product into the nostril of children. A small amount applied this way may cause immediate collapse.

Interactions with Other Medicaments:

Warfarin- Excess application on the skin may increase the chances for bleeding.

Side Effects:

Skin irritation, redness or swelling, burning or stinging sensation.
Adverse reaction is rare, symptom of dizziness, nausea may occur.

Symptoms and Treatment of Overdose:

Ingestion may cause gastrointestinal disturbances such as nausea, dyspepsia and vomiting. Other irritation of the gastric mucosa with ulceration, erosion, haematemesis, melaena and slight blood loss may occur.

Symptoms of poisoning includes nausea, vomiting, acidosis, pulmonary edema, pneumonia, convulsions and death.

Treatment:

The stomach should be emptied by gastric lavage. Salicylate remaining in the stomach may be absorbed by activated charcoal. Fluid and electrolyte management is the mainstay of the treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalaemia and dehydration. Alkaline diuresis, haemodialysis or haemoperfusion are effective methods of removing salicylate from the plasma.

Storage Conditions:

Store below 30°C, in a dry place, protected from direct sunlight.

Dosage Forms:

Ointment

Pack Size:

30gm and 60gm HDPE tub and Alu tube.
450gm HDPE jar (for tender purpose).

Registration Number:

MAL06021262X

Manufacturer by:

KCK PHARMACEUTICAL INDUSTRIES SDN. BHD.
Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Product Registration Holder:

UNIMED SDN. BHD.
53, Jalan Tembaga SD 5/2B,
Bandar Sri Damansara,
52200 Kuala Lumpur, Malaysia.

Date of Revision:

11 April 2023