

110mm

UNI-PAIN LINIMENT

Name and Strength of Active Substance:

Each 100 ml contains:

Methyl Salicylate	25 gm
Preservative:	
Methyl Paraben	0.1% w/v
Contains Alcohol	2.0% v/v

Product Description:

Yellow colour liniment with a characteristic odour.

Indication:

For temporary relief of minor muscular and rheumatic pains.

Application:

Apply a little to the affected part and gently massage. To be applied 3 or 4 times a day.

Contraindications:

Intolerance of one of the component of the preparation. This product is contraindicated in child below 2 years of age. Caution should be exercised when older children are treated..

Warning and Precautions:

- For external use only.
- If condition worsen, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.
- Do not apply to wound or damaged skin.
- Do not apply over large areas of the body or bandage.
- This product contains methyl salicylate and when applied or rub to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application of the skin for muscle or joint pain may increase the chances of bleeding.

Interactions with Other Medicaments:

Warfarin - Excess application on the skin may increase the chances for bleeding.

Side Effects:

Skin irritation, redness or swelling, burning or stinging sensation.
Adverse reaction is rare, symptom of dizziness, nausea may occur.

Symptoms and Treatment of Overdose:

Ingestion may cause gastrointestinal disturbances such as nausea, dyspepsia and vomiting. Other irritation of the gastric mucosa with ulceration, erosion, haematemesis, melaena and slight blood loss may occur.

Treatment:

The stomach should be emptied by gastric lavage. Salicylate remaining in the stomach may be absorbed by activated charcoal. Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalaemia and dehydration. Alkaline diuresis, haemodialysis or haemoperfusion are effective methods of removing salicylate from the plasma.

Storage Conditions:

Store below 30°C, in a dry place, protected from direct sunlight.

Dosage Forms:

Liniment

Pack Size:

60ml, 90ml, 100ml, 120ml, 200ml and 250ml in HDPE bottle.

Registration Number:

MAL06021263XC

Manufactured by:

KCK PHARMACEUTICAL INDUSTRIES SDN. BHD.
No 5 & 7 Hala Beringin,
Diamond Valley Industrial Park,
11960 Bayan Lepas, Pulau Pinang, Malaysia

Product Registration Holder:

UNIMED SDN. BHD.
53, Jalan Tembaga SD 5/2B,
Bandar Sri Damansara,
52200 Kuala Lumpur, Malaysia.

Date of Revision:

24 June 2024

PI0038-1

193mm