

PABRON™ COUGH SYRUP 100mg/5ml kids

パブロン



- Non-Drowsy
- Loosens & Clears Phlegm
- Relieves Cough
- Alcohol-Free & Sugar-Free
- Blackcurrant-Flavoured



PABRON™
COUGH SYRUP
100mg/5ml
kids



PABRON™
COUGH SYRUP
100mg/5ml
kids



PABRON™
COUGH SYRUP
100mg/5ml
kids



Each 5 ml syrup contains:
Carbocisteine 100mg
Methyl Hydroxybenzoate
as Preservative 0.15% w/v

Product Description

A reddish purple syrup with blackcurrant flavour.

Pharmacodynamics

Carbocisteine is a mucolytic. It alters the structure of the mucus and decreases its viscosity in order to facilitate its removal by ciliary action or expectoration.

Pharmacokinetics

Carbocisteine is rapidly and well absorbed from the gastrointestinal tract. It appears to penetrate lung tissue and respiratory mucus.

Carbocisteine is excreted in the urine as unchanged drugs and metabolites.

Acetylation, decarboxylation and sulfoxidation have been identified as the major metabolic pathways.

Indications

Pabron Kids Cough aids expectoration by reducing sputum viscosity, and therefore loosens and clears phlegm and relieves cough.

Recommended Dosage

Children above 5 years:
5ml 3 times daily
Children from 2 to 5 years:
2.5ml 3 times daily

Side Effects

Nausea, gastric discomfort, gastrointestinal bleeding and skin rash have occurred occasionally.

Immune System Disorders:

Anaphylactic/ anaphylactoid reaction.

Skin and Subcutaneous Tissue Disorders:

Severe cutaneous adverse reactions (SCAR) e.g. erythema multiforme, Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN). In most of these cases reported at least one other drug administered at the same time, which may have possibly enhanced the described mucocutaneous effects.

Contraindications

Patient with active peptic ulcers.
Contraindicated in children under 2 years of age.

Warning & Precautions

During treatment, do not take any anti-cough medicine or any drug which dries bronchial secretion. Inform your doctor if you are pregnant or breastfeeding. If in doubt, do not hesitate to contact your doctor or pharmacist.

Interactions with Other Medication

In order to avoid possible interactions when taking more than one drug, all other treatments you are taking must be routinely reported to your doctor or pharmacist.

Symptoms & Treatment for Overdose

In the event of overdose, stop the treatment and consult your doctor as soon as possible.

Pregnancy & Lactation

Safe use of Carbocisteine during pregnancy and lactation has not been established.

Route of Administration: Oral

Packing:

120ml in amber PET bottle.

Storage

Store below 30°C and away from heat.

KEEP OUT OF REACH OF CHILDREN

For further information, please consult your pharmacist and physician.

Shelf-Life

2 years from the date of manufacture.

Manufactured by:

Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate, 75250
Melaka, Malaysia

Under Licence of:

Taisho Pharmaceutical Co., Ltd. Tokyo,
Japan

Product Registration Holder:

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6,
Bandar Sultan Suleiman,
42000 Port Klang,
Malaysia.

Revision date: May 2018

PA0500-003

