

RANCANGAN BROSUR BALSEM GOSOK - BALSEM MERAH

CAP BETET 7,5 g dan 17,5 g EM

SISI DEPAN

SISI BELAKANG



BALSEM GOSOK - BALSEM MERAH CAP BETET

Composition/Kandungan :

Methyl Salicylate

10% w/w

Camphor

10% w/w

Menthol

9% w/w

Bentuk Dosej : Salep.

Desripsi Produk : Salep yang berwarna merah dan berbau mentol.

Farmakodinamik : Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakokinetik : Sila rujuk kepada penerangan Bahasa Inggeris.

Indikasi : Untuk melegakan sakit otot, sakit sendi, sakit urat dan terseliuh.

Arahan Penggunaan : Sapu ubat ini pada bahagian berkenaan 3 kali sehari. Jangan balut bahagian tersebut.

Cara Penggunaan : Topikal.

Kontraindikasi : Produk ini tidak sesuai untuk pesakit yang hipersensitif kepada bahan aktif.

Amaran dan Perhatian : Produk ini boleh menyebabkan sawan, tidak sesuai untuk bayi bawah umur 2 tahun. Perhatian diberikan kepada kanak-kanak berumur 2 tahun ke atas bila menggunakannya. Sapu ubat yang mengandungi camphor pada hidung kanak-kanak adalah merbahaya kerana boleh menyebabkan pengsan.

Produk ini mengandungi methyl salicylate dan ia mungkin boleh diserap masuk ke dalam darah melalui kulit. Penyapuan yang berlebihan oleh pesakit yang mengambil warfarin mungkin meningkatkan risiko pendarahan.

Tidak boleh disapu pada bahagian mata, permukaan mukus, luka terbuka atau kulit yang rosak. Elakkan penggunaan yang berlebihan semasa mengandung dan penyusuan. Jangan sapu secara meluas pada bahagian badan atau membalut. Jika iritasi kulit berlaku, sila berhenti guna.

Interaksi dengan ubat lain : Warfarin.

Kehamilan dan Penyusuan : Keselamatan penggunaan oleh wanita hamil tidak dapat dipastikan. Oleh itu produk ini harus digunakan dengan berhati-hati supaya tidak mandatkan risiko kepada ibu dan anak.

Kesan Sampingan : Iritasi kulit, kemerahan, bengkak, rasa pedih berlaku. Reaksi buruk jarang berlaku, simptom pening, loya boleh berlaku.

Symptom dan Rawatan Overdos : Sila rujuk kepada penerangan Bahasa Inggeris.

Dosage Form : Ointment.

Product Description : A red colour ointment with menthol smell.

Pharmacodynamics

Camphor

is a stimulant, rubefacient, antipruritic and feebly antiseptic. When in combination with other ingredient like menthol, methyl salicylate, it becomes ideal for neuralgia and other painful areas.

Menthol

dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Methyl salicylate

is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins, and cause platelet dysfunction. Methyl salicylate is used topically as a counter-irritant. Upon application, it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscle or joints.

Pharmacokinetics

Camphor

is readily absorbed from all administration sites. Camphor is well absorbed after dermal exposure. It is hydroxylated in the liver to yield hydroxycamphor metabolites which are then conjugated with glucuronic acid and excreted in the urine. Camphor crosses the placenta.

Menthol

After absorption, menthol is excreted in the urine and bile as a glucuronide.

Methyl salicylate

May be absorbed through intact skin. The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity. Absorption characteristics of salicylates vary with the dose, formulation and route of administration. Percutaneous absorption is enhanced by exercise, heat, occlusion, or disruption of the integrity of the skin or application to large areas of skin. Both the rate and extent of absorption increases after repeated application; increasing the bioavailability. Methyl salicylate is extensively metabolised to salicylic acid in the dermal and subcutaneous tissues after topical application. At therapeutic levels, the half life of salicylates is 2-4 hours. As salicylate level increase into the toxic range, the half-life can be greater than 18 hours.

Indication : For the relief of muscle pain, joint pain and sprain.

Direction for use : Apply on the affected area, 3 times daily. Do not bandage.

Route of Administration : Topical application.

Contraindication : Patient with known hypersensitivity to the active ingredients.

Warning and Precaution : This product can cause convulsion and is contraindicated in infants below 2 years of age. Caution must be exercised when older children are treated. It is dangerous to place any camphor containing product into the nostril of children. A small amount applied this way may cause immediate collapse.

This product contains methyl salicylate and when applied or rub on the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

Avoid contact with the eyes and mucous membrane. Do not apply to open wounds or damaged skin. Avoid excessive use during pregnancy and lactation. Do not apply over large areas of the body or bandage. Discontinue use if irritation develops.

Drug Interaction : Warfarin.

Pregnancy and Lactation : Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child.

Side Effects : Skin irritation, redness or swelling, burning or stinging sensation. Adverse reaction is rare, symptom of dizziness, nausea may occur.

Symptoms and Treatment of Overdose : Mild chronic salicylate intoxication, or salicylism, usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache, and confusion may be controlled by reducing the dosage.

Storage/Penyimpanan : Store below 30°C. Keep out of reach of children. For external use only. Not to be taken.

Simpan di bawah 30°C. Jauhkan daripada kanak-kanak. Untuk kegunaan luar sahaja. Tidak boleh dimakan.

Packing/Bungkusan : 7.5 g and 17.5 g

Shelf Life/Tempoh Hayat : 5 years

Date of revision : January 2025

Dikilang oleh Pemilik Produk : PT IRAWAN DJAJA AGUNG

Jalan Raya Sukodono, Kecamatan Sukodono, Sidoarjo, Jatim, Indonesia.

Pemegang Produk, Pengimport dan Pengekar : Pasir Mas-Indo Trading Sdn Bhd.

No.52 (GF), Lorong Kelasah 3, Taman Kelasah, 13700 Perai, Pulau Pinang, Malaysia.