

Live-well™ OSTESAMIN 500 Capsule

COMPOSITION:

Each vegetable capsule contains:

Glucosamine sulphate sodium chloride 628mg equivalent to Glucosamine Sulphate 500mg.

PRODUCT DESCRIPTION:

Two pieces, yellow colored hard vegetable capsule of size “00” containing white to light brown powder.

PHARMACOLOGY:

Glucosamine stimulates the biosynthesis of mucopolysaccharides – the key structural components of joint cartilage which are essential for proper joint function. Glucosamine also increases synovial fluid production and viscosity, giving better lubricant activity for the joints.

Glucosamine is effective in reducing articular pains, improving articular function and inhibiting or slowing down the process of degenerative osteoarticular diseases.

Glucosamine sulphate is well-absorbed after oral intake, and is distributed mainly to bones and joint tissues.

PHARMACOKINETICS:

After oral administration, 90% of glucosamine sulphate is absorbed in the small intestine and then transported to the liver via portal circulation. Free glucosamine is not detected in the serum. In animal studies, glucosamin is shown to diffuse into several tissues, including bones, and articular cartilages.

INDICATION:

As adjuvant therapy for osteoarthritis.

RECOMMENDED DOSE:

Light or moderate osteoarthritic symptoms: Take one capsule twice daily for at least six weeks.

Severe osteoarthritic symptoms: Start with one capsule three times daily for at least 8 weeks. Then take one capsule twice daily for another 3-4 months (or as professionally prescribed).

For long-term therapy, treatment should be repeated every other six months or at shorter intervals, as professionally prescribed.

Capsules should be taken 15 minutes before meals.

Glucosamine sulphate treats the cause of osteoarthritis and pain relief can be seen only after about one week. Therefore, if severe pain is present during this period, it may be necessary to take an analgesic such as aspirin or a non-steroidal anti-inflammatory drug (NSAID).

CONTRAINDICATIONS:

Contraindicated in individuals who are hypersensitivity to glucosamine.

WARNING AND PRECAUTION:

Derived from seafood.

The effect of glucosamine on insulin sensitivity and glucose tolerance is unclear, caution is advised for diabetic patients.

INTERACTIONS WITH OTHER MEDICATIONS:

None known

PREGNANCY AND LACTATION

If you are pregnant or breastfeeding, please consult your doctor or health care professional before taking this product.

SIDE EFFECTS:

Cardiovascular

Peripheral oedema, tachycardia were reported in few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

Central Nervous System

Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%)

Gastrointestinal

Nausea, vomiting, diarrhea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine.

Skin

Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

OVERDOSE AND TREATMENT

In the event of an acute overdose, the patient should be observed closely and symptomatic treatment should be instituted.

STORAGE CONDITION:

Keep bottle tightly closed.

Store in a dry place below 30°C.

SHELF-LIFE

3 Years

PACK SIZE:

7, 10, 30, 60, 90 capsules per bottle.

Blister of 10 capsules. Box of 100, 500, 1000 capsules.

MANUFACTURED BY/FOR:

Pahang Pharmacy Sdn. Bhd. (10826-K)

Lot 5979, Jalan Teratai, 5 ½ Miles Off Jalan Meru,
41050 Klang, Selangor Darul Ehsan.

Revised date: 15 October 2008