

5/25/2022
17630 MY Procosa II

Label size: 2.5 x 6 150cc PB

Colors:
7544 C
644 C
646 C
647 C
Black

USANA


procosa[™] II

84 Tablets
MAL06061538X



Directions: Take one (1) tablet three times per day, preferably with meals. Each tablet contains glucosamine sulphate equivalent to glucosamine 300 mg. Keep out of reach of children. Jauhkan dari kanak-kanak.

Serving Size: 1 Tablet	Amount Per Serving	%DV*
Glucosamine	300 mg	†

*Daily Value based on U.S. recommended daily allowance.
†Daily Value not established.

Other Ingredients: Microcrystalline Cellulose, Ascorbyl Palmitate, Dextrin, Colloidal Silicon Dioxide, Dextrose, Soy Lecithin, Sodium CMC, Sodium Citrate.
Pregnancy and breastfeeding: Insufficient reliable data.
There is a safety seal under the cap. Do not use if the safety seal is broken or missing.
Derived from seafood.
Please refer to enclosed leaflet for details.
Production Registration Holder: / Imported by: UHS Essential Health Malaysia Sdn. Bhd.
Unit M2-2 & M2-5, Level M2, The Vertical Podium, Avenue 3, Bangsar South, No. 8, Jalan Kerinchi, 59200 Kuala Lumpur, Malaysia
Manufactured by: USANA Health Sciences, Inc., 3838 W. Parkway Blvd., Salt Lake City, Utah 84120 USA
Store in a dry place at or below 30 °C. Protect from light and moisture.
131.010104 LB.002424 Made in USA.



DESCRIPTION

White, modified rectangular tablet, clear film-coated with no imprinting. Some colour variation is normal.

PHARMACODYNAMICS

Biochemically, glucosamine is involved in glycoprotein metabolism. Glycoproteins, known as proteoglycans, form the ground substance in the extra-cellular matrix of connective tissue. Proteoglycans are polyanionic substances of high-molecular weight and contain many different types of heteropolysaccharide side-chains covalently linked to a polypeptide-chain backbone. These polysaccharides make up to 95 per cent of the proteoglycan structure. In fact, chemically proteoglycans resemble polysaccharides more than they do proteins.

The polysaccharide groups in proteoglycans are called glycosaminoglycans or GAGs. GAGs include hyaluronic acid, chondroitin sulphate, dermatan sulphate, keratan sulphate, heparin, and heparan sulphate. All of the GAGs contain derivatives of glucosamine or galactosamine.

Glucosamine derivatives are found in hyaluronic acid, keratan sulphate, and heparan sulphate. Chondroitin sulphate contains derivatives of galactosamine.

The glucosamine-containing glycosaminoglycan hyaluronic acid is vital for the function of articular cartilage. GAG chains are fundamental components of aggrecan found in articular cartilage. Aggrecan confers upon articular cartilage shock-absorbing properties. It does this by providing cartilage with a swelling pressure that is restrained by the tensile forces of collagen fibres. This balance confers upon articular cartilage, the deformable resilience vital to its function.

In the early stages of degenerative joint disease, aggrecan biosynthesis is increased. However, in later stages, aggrecan synthesis is decreased, leading eventually to the loss of cartilage resiliency and to most of the symptoms that accompany osteoarthritis.

During the progression of osteoarthritis, exogenous glucosamine may have a beneficial role. It is known that, in vitro, chondrocytes do synthesise more aggrecan when the culture medium is supplemented with glucosamine. N-acetylglucosamine is found to be less effective in these in vitro studies. Glucosamine has also been found to have antioxidant activity and to be beneficial in animal models of experimental arthritis.

The counter anion of the glucosamine salt (i.e., chloride or sulphate) is unlikely to play any role in the action or pharmacokinetics of glucosamine. Further, the sulphate in glucosamine sulphate preparations should not be confused with the glucosamine sulphate found in such GAGs as keratan sulphate and heparan sulphate. In the case of the preparation, sulphate is the anion of the salt. In the case of the above GAGs, sulphate is present as an ester. Also, there is no glucosamine sulphate in chondroitin sulphate.

PHARMACOKINETICS

Pharmacokinetics of glucosamine are derived primarily from animal studies. About 90 per cent of glucosamine administered orally as glucosamine salt gets absorbed from the small intestine, and from there it is transported via the portal circulation to the liver. It appears that a significant fraction of the ingested glucosamine is catabolised by first-pass metabolism in the liver. Elimination of glucosamine is primarily through the urine, with a small amount of glucosamine or its derivatives eliminated in the faeces.

INDICATION

As an adjunctive for the treatment of osteoarthritis.

RECOMMENDED DOSAGE

Take one (1) tablet three times per day, preferably with meals. Each tablet contains glucosamine sulphate 2 KCl, equivalent to glucosamine 300 mg.

MODE OF ADMINISTRATION

Oral.

CONTRAINDICATIONS

Patients who are hypersensitive to glucosamine. Patients who are allergic to shellfish/seafood.

WARNINGS AND PRECAUTIONS

Glucosamine increases insulin resistance in normal and experimentally diabetic animals. In these animals, intravenous glucosamine significantly decreases the rate of glucose uptake in skeletal muscle. In animals given oral glucosamine, this is not observed.

Those with Type 2 diabetes and those who are overweight and have problems with glucose tolerance should have their blood sugar carefully monitored if they use glucosamine preparations.

Glucosamine is not meant to replace nsaids in cases of intense pain, as this is a supportive therapy.

This product is derived from seafood.

INTERACTIONS WITH OTHER MEDICAMENTS

Diabetics who, under medical advisement, decide to use glucosamine preparations will need to monitor their blood glucose and may need to adjust the doses of the medications they take to control blood glucose. This needs to be done under medical supervision. No other drug, nutritional supplement, food, or herb interaction is known.

PREGNANCY AND LACTATION

Because of insufficient safety data, children, pregnant women, and nursing mothers should avoid using glucosamine.

SIDE EFFECTS

Side effects of glucosamine that have been reported are mainly mild gastrointestinal complaints such as heartburn, epigastric distress, and diarrhoea.

- Cardiovascular: Peripheral oedema and tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.
- Central nervous system: Drowsiness, headache, and insomnia have been observed rarely during therapy (less than 1%).
- Gastrointestinal: Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation, heartburn, and anorexia have been described rarely during oral therapy with glucosamine.
- Skin: Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

SYMPTOMS AND TREATMENT OF OVERDOSE

None known.

HOW SUPPLIED

84 tablets in a plastic bottle.

STORAGE CONDITION

Store below 30 °C. Protect from light and moisture.

SHELF LIFE

24 months.

MANUFACTURED BY:

USANA Health Sciences, Inc.
3838 W. Parkway Blvd., Salt Lake City, Utah 84120 USA

PRODUCT REGISTRATION HOLDER / IMPORTED BY:

UHS Essential Health (Malaysia) Sdn. Bhd.
Unit M2-2 & M2-5, Level M2, The Vertical Podium, Avenue 3, Bangsar South, No. 8, Jalan Kerinchi, 59200 Kuala Lumpur, Malaysia

LOT - MFG - EXP
has been added. I have moved
the Halal and recycle symbol to
make more room for this if it
is laser printed. Please let me
know if this is placed in a good
spot for this and other labels
going forward. Thanks.