



DANZEN® TABLETS
Antiinflammatory & antitumefacient
oral enzyme preparation

COMPOSITION:-
Each tablet contains 5 mg serratiopeptidase (10,000 serratiopeptidase units).

DESCRIPTION:-
Light orange, enteric film coated tablet, printed logo " Ⓢ " in black ink on one face. The diameter is 7.3 mm.

PHARMACODYNAMICS & PHARMACOKINETICS:-
1) Anti-inflammatory and Anti-swelling actions.
a. Serratiopeptidase, orally given, represses the fibrinolysis activity and increases vascular permeability in the scalded rat, as an inflammation model.
b. Serratiopeptidase, orally given, inhibits inflammatory edema induced by anti-rat serum in the rat. Similarly, the drug inhibits inflammatory edema in the rat due to various kinds of phlogistic agents, including carrageenin, dextran, serotonin and bradykinin.
c. Serratiopeptidase strongly hydrolyzes bradykinin, a phlogistic polypeptide (in vitro).
d. Serratiopeptidase strongly lyzes fibrin and fibrinogen, but does not appreciably affect proteins in the living body, such as albumin and α- and γ-globulins.
2) Enhancing effect on the lysis and discharge of sputum and pus.
a. Serratiopeptidase appears to decrease the dry weight and viscosity of nasal discharge in patients with chronic sinusitis.
b. In rabbits with subacute bronchitis, induced by exposure to sulfurous acid gas, serratiopeptidase decreases the dry weight of the sputum, resulting in a significant decrease in the sputum viscosity. It was also confirmed that the serratiopeptidase decreased the dry weight of nasal discharge in patients with chronic sinusitis.
3) Enhancing the effect on the transfer of antibiotics to the focal site.
Serratiopeptidase accelerates the transfer of ampicillin and sulbenicillin to the maxillary sinus mucosa in patients with chronic sinusitis. The drug also accelerates the transfer of mecillinam into the bladder wall of rabbits with experimental cystitis.

Although a couple of studies on absorption of serratiopeptidase have already reported, Miyata et al. confirmed by using the radioimmunoassay (RIA) in rats and dogs that serratiopeptidase itself was absorbed through the intestinal tract.
Further, Yashiki reported by the RIA method that serratiopeptidase was absorbed through the intestinal tract after oral administration of 5 mg enteric coated tablets. It was also confirmed that serratiopeptidase was bound to alpha-2 macroglobulin to form the complexity in the body and that serratiopeptidase in the blood was quickly transferred to the focal site (rats). Nakahama et al. elucidated by using biotechnological methods the complete amino acid sequence (470 amino acids) and, consequently, the molecular weight (50,632) of serratiopeptidase.

- INDICATIONS:-**
(1) Inflammation in the following diseases:
1) After surgery and trauma
2) In the field of otorhinolaryngology: Sinusitis
3) In the field of obstetrics & gynecology: Engorgement of breast
4) In the field of urology: Cystitis and epididymitis
5) In the field of dental & oral surgery: Pericoronitis of the wisdom tooth and alveolar abscess
(2) Difficulty of expectoration in the following diseases:
Bronchitis, pulmonary tuberculosis and bronchial asthma
(3) Difficulty of expectoration after anesthesia

DOSAGE AND ADMINISTRATION:-
Usually, take one or two tablets of *DANZEN®* orally, three times a day after meals. Swallow without chewing, together with a glass of water or lukewarm water. The dosage should be adjusted according to the patient's age and conditions.

143400016
P3/19 MS E1/2

CONTRAINDICATIONS:-
Patients with a history of hypersensitivity to the ingredients of this drug.

- WARNINGS AND PRECAUTIONS:-**
- Indications and Dosage and Administration should be adhered to.
 - *DANZEN® Tablet* will not interfere with the capability of driving a car or operating heavy machinery.
 - Consultation with the doctor or pharmacist is needed in case of the following patients:
1) Patients with a history of drug allergy
2) Patients with blood coagulation abnormalities, such as epistaxis and bloody sputum; severe hepatic or renal dysfunction or under treatment with anticoagulants.
 - The mechanism of action of *DANZEN®* in the body is not fully clarified and a definite dose-response relationship has not been established as yet. Therefore, it should not be administered desultorily.

INTERACTIONS WITH OTHER MEDICAMENTS:-
Since the concomitant use of *DANZEN®* with an anticoagulant may intensify the anticoagulant effect, *DANZEN®* should be administered cautiously under close observation.

PREGNANCY AND LACTATION:-
No data available

ADVERSE EFFECTS / UNDESIRABLE EFFECTS:-
The following symptoms may occur. If such symptoms occur, *DANZEN® Tablet* should be discontinued and consult with your doctor or pharmacist.
- Skin: Stevens-Johnson syndrome, Toxic epidermal necrolysis, Rash or redness, Pruritus
- Hypersensitivity: Shock, Anaphylactic symptoms
- Hepatic: Hepatitis, Jaundice
- Gastrointestinal: Diarrhea, anorexia, gastric discomfort, nausea or vomiting
- Others: Epistaxis or bloody sputum

OVERDOSE:-
There is no specific data to be described.

STORAGE CONDITIONS:-
Store below 30°C, away from humidity and light.

SHELF LIFE: -
4 years

DOSAGE FORMS AND PACKAGING AVAILABLE:-
100 tablets (10 tablets / aluminium blister sheet) in a box
120 tablets (10 tablets / aluminium blister sheet) in a box

Manufactured by
INTERTHAI PHARMACEUTICAL MANUFACTURING LIMITED,
1899 Phaholyothin Road, Ladyao, Chatuchak, Bangkok, Thailand

For
LUEN WAH MEDICAL CO (S) PTE LTD, SINGAPORE

Product Registration Holder:-
LUEN WAH MEDICAL CO. SDN. BHD.
No. 53-2, 49-G, 49-1, The Highway Centre, Jalan 51/205, 46050 Petaling Jaya, Selangor Darul Ehsan, Malaysia

DATE OF REVISION OF INSERT:- 29 Jan 2019

143400016
P3/19 MS E2/2

Client : Interthai	Date : 28/11/2019	References Color : 1 Color Black		Client
Product : Leaflet Danzen Tablets (MS)				Approved by :
Code : 143400016 P3/19	Dimension : 120x180 mm			Date :

TEL. 0-2840-7117 FAX. 0-2840-7118 www.ulprint.co.th