

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory.

IOBET EYE DROPS

Betaxolol Ophthalmic Solution USP

DISCRIPTION:

Clear, Colourless solution

NAME(S) AND STRENGTH (S) OF ACTIVE INGREDIENTS(S):

Betaxolol0.5% w/v

(As equivalent to Betaxolol hydrochloride USP0.56% w/v)

Benzalkonium chloride NF as preservative0.01% w/v.

DOSAGE FORM: Eye Drops

PHARMACODYNAMICS:

Ophthalmology: Antiglaucoma Preparations & Miotics. Betaxolol is a cardioselective Beta receptor blocker which, when applied topically to the eye, lowers intraocular pressure. It is thought to produce this effect by reducing the rate of production of aqueous humour.

Clinical Pharmacology : Several studies have indicated that betaxolol may have a beneficial effect on visual function for up to 48 months in patients with chronic open- angle glaucoma and up to 60 months in patients with ocular hypertension. Moreover there is evidence that betaxolol maintains or increase ocular blood flow/ perfusion.

PHARMACOKINETICS:

Betaxolol is highly lipophilic which results in good permeation of the cornea, allowing high intraocular levels of the drug. Betaxolol is characterized by its good oral absorption, low first pass loss and a relatively long half-life of approx 16-22 hours. The elimination of betaxolol is primarily by the renal rather than faecal route. The major metabolic pathways yield two carboxylic acid forms plus unchanged betaxolol in the urine(approx. 16% of the administered dose).

INDICATION

Iobet is indicated for the reduction of elevated intraocular pressure in patients with ocular hypertension and chronic open angle glaucoma.

CONTRA INDICATIONS

Hypersensitivity to any component of this product. Betaxolol 0.5% eye drops are contraindicated in patients with sinus bradycardia, greater than a first degree atrioventricular block, cardiogenic shock, or patients with overt cardiac failure.

DOSAGE AND ADMINISTRATION:

Children and Adults(including the elderly) the usual dose is one drop to be instilled into the affected eye(s) twice daily. Children : Iobet is not recommended for use in children. Use the solution within one month after opening the vial.

USE IN PREGNANCY AND LACTATION:

There have been no adequate and well controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response. This drug should be used during pregnancy only if clearly indicated. Beta-adrenergic receptor blocking agents may cause bradycardia in the foetus and new born infant. During the final part of the pregnancy and parturition these drugs should, therefore, only be given after weighing the needs of the mother against the risk to the foetus. It is not known whether ophthalmic betaxolol hydrochloride is excreted in human milk., therefore caution should be exercised when ophthalmic betaxolol hydrochloride is prescribed for breast-feeding women.

SYMPTOMS AND TREATMENT OF OVERDOSE:

A topical overdose of Iobet may be flushed from the eye(s) with warm tap water.

WARNING:

Patients who are receiving a beta-adrenergic blocking agent orally and Iobet should be observed for potential additive effect either on intraocular pressure or the known systemic effects of beta-blockade. While Iobet has demonstrated a low potential for systemic effects, it should be used with caution in patients with diabetes(especially labile diabetes) or in patients suspected of developing thyrotoxicosis. Consideration should be given to the gradual withdrawal of beta-adrenergic blocking agents prior to general anaesthesia because of the reduced ability of the heart to respond to beta-adrenergically mediated sympathetic reflex stimuli. Iobet, a cardioselective beta-blocker, has produced only minimal effects in patients with reversible airway obstruction, however, caution should be exercised in the treatment of patients with a history of obstructive pulmonary disease. In patients with angle-closure glaucoma, the immediate treatment objective is to re-open the angle by constriction of the pupil with a miotic agent, betaxolol has no effect on the pupil, therefore, Iobet should be used with a miotic to reduce elevated intraocular pressure in angle-closure glaucoma. This product contains benzalkonium chloride and is not recommended for use when soft contact lenses are being worn.

SIDE EFFECTS:

Although Iobet is generally well tolerated, discomfort of short duration may be experienced by some patients upon instillation and occasional tearing has been reported.

Ocular Effects

Common: discomfort

Uncommon: tearing Rare: decreased corneal sensitivity, erythema, itching, corneal punctate staining, keratitis, anisocoria and photophobia.

Systemic reactions following topical administration of betaxolol have been rarely reported.

Systemic Effects:

Nervous System : Rare: insomnia, depression

Body as a whole : Rare: headache

Cardiovascular: Rare: bradycardia

Respiratory: Rare: dyspnoea, asthma.

Skin and Appendages: Rare: alopecia

Since topically applied beta-adrenergic blocking agents may be absorbed systemically, adverse reactions found with systemic administration of beta adrenergic blocking agents may occur with topical administration. These may include bradycardia, a slowed AV conduction or increase of an existing AV block, hypotension, heart failure, cold and cyanotic extremities, Raynaud phenomenon, paraesthesia of the extremities, increase of an existing intermittent claudication, fatigue, headaches, impaired vision, hallucinations, psychoses, confusion, impotence, dizziness, sleep disturbances, depression, nightmares, gastro-intestinal problems, nausea, vomiting, diarrhoea, bronchospasm in patients with bronchial asthma or history of asthmatic complaints, disorder of the skin, especially rash, and dry eyes. Beta blockers may mask the symptoms of thyrotoxicosis or hypoglycemia. An increase in Anti Nuclear Antibodies (ANA) has been seen ; its clinical relevance is unclear.

DRUG INTERACTIONS:

Patients who are receiving a beta-adrenergic blocking agent orally and should be observed for a potential additive effect either on the intraocular pressure or on the known systemic effects of beta blockade. Close observation of the patient is recommended when a beta blocker is administered to patients receiving catecholamine-depleting drugs such as reserpine, because of possible additive effects and the production of hypotension and/or bradycardia. Betaxolol is an adrenergic blocking agent; therefore caution should be exercised in patients using concomitant adrenergic psychotropic drugs.

STORAGE CONDITION:

Store the vial well closed at a temperature below 30°C. Discard container one month after opening.

SHELF LIFE:

24 Months from the date of Manufacturing

THERAPEUTIC CLASS:

Anti Glaucoma Agent.

Presentation:

Iobet Eye drops available in 5 ml of plastic container

MODE OF USE




SPIKE

Tighten the cap on the nozzle



The spike in the cap will pierce the tip of the vial



Dispense drops with gentle pressure

Replace the cap after every use

Imported By:
Unimed SDN BHD
No. 53, Jalan Tembaga SD 5/2B.
Bandar Sri Damansara 52200,
Kuala Lumpur, Malaysia

Manufactured in India by:

FDC FDC Limited

B-8, MIDC Industrial Area, Waluj,
Aurangabad - 431 136, Maharashtra.

W1BLL01AESMY

EXPORT	RA	QA	R&D (F)	Packaging Development Department	Checked by	Approved by

HISTORY: 1) Artwork revised due to old colde.
2) Pharmacode position changed.

Mini Pharma Code No. : Front: 720/Back: 725		FDC Limited LEAFLET	
Size	110(L)x144(H) mm	Unfolded Size	
Folded Size		No. of Folds	
GSM	60 gsm	Paper	Maplitho Paper
Specifcan No.		Colour	TEXT IN BLACK
Artwork Code: W1BLL01AESMY	Supercodes AW Code: EXG-WI-0021	Location : Waluj - Malaysia	Prepared On : 17-04-14