

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

Cycloserine Capsules USP 250 mg

Coxerin

COMPOSITION

Each capsule contains:

Cycloserine USP.....250 mg

EXCIPIENTS

Lightmagnesium oxide BP, Heavy magnesium oxide BP, Purified Talc BP, Maroon/Maroon EHG, Size '1'

DESCRIPTION

Maroon/Maroon Size "1" hard gelatin filled capsule filled with white to pale yellow powder.

ACTION

Cycloserine inhibits cell wall synthesis in susceptible organisms by competing with D alanine for incorporation into the bacterial cell wall.

CLINICAL PHARMACOLOGY

Cycloserine is readily absorbed from the gastrointestinal tract, with peak blood levels occurring in 4 to 8 hours. Concentration in the CSF, pleural fluid, fetal blood, and mother's milk approach those found in serum. Detectable amounts are found in ascetic fluid, bile, sputum, amniotic fluid, lung tissues and lymph tissues. Approximately 65% of a single dose of cycloserine can be recovered in the urine within 72 hours after oral administration. The remaining 35% is apparently metabolized to unknown substances. The maximum excretion rate occurs 2 to 6 hours after administration, with 50% of the drug eliminated in 12 hours.

INDICATIONS

Cycloserine is indicated in the treatment of multidrug resistance tuberculosis in combination with other anti-tubercular drugs.

DOSAGE

In the treatment of tuberculosis and other mycobacterial disease, cycloserine should not be given alone and therapy should be continued long enough to prevent relapse. In adults with multidrug resistant tuberculosis cycloserine therapy is usually initiated with 250 mg every 12 hours. Dosage should be reduced in patients with renal impairment and if possible, serum concentrations monitored weekly. The dosage should be adjusted to keep the blood levels below 30mcg/ml.

SIDE EFFECTS

The most frequent adverse effects of cycloserine involve the nervous system. Drowsiness, dizziness, headache, lethargy, depression, **psychosis**, tremor, dysarthria, nervousness, anxiety, vertigo, confusion and major and minor chronic seizures, convulsions and coma have been reported in patients receiving the drug.

Adverse nervous system effects appear to be dose related and occur within the first 2 weeks of therapy in about 30% of patients receiving 500 mg of cycloserine daily; symptoms generally disappear when the drug is discontinued. Rarely, hypersensitivity reactions including rash and photo-sensitivity have occurred with cycloserine. Cardiac arrhythmias and sudden development of congestive heart failure have been reported rarely in patients receiving 1 - 1.5 gm of the drug daily.

CONTRAINDICATIONS

Cycloserine is contraindicated in patients with a history of hypersensitivity, epilepsy, mental depression, severe anxiety or psychosis. The drug is also contraindicated in patients with severe renal impairment, in frequent users of alcohol.

DRUG INTERACTIONS

Ethionamide, Isoniazid - concurrent administration potentiates neurotoxicity.

Alcohol - increases the risk of epileptic episodes

SYMPTOMS AND TREATMENT OF OVERDOSE

Patients receiving more than 500 mg daily should be closely observed for the signs of CNS toxicity - convulsions, psychosis, somnolence, depression, confusion, hyperreflexia, headache, tremor, vertigo, paresis.

WARNINGS

Administration of Coxerin should be discontinued or the dosage reduced if the patient develops allergic dermatitis or

symptoms of CNS toxicity, such as convulsions, psychosis, somnolence, depression, confusion, hyperreflexia, headache, tremor, vertigo, paresis,

PRECAUTIONS

Before initiation of treatment, the organism's susceptibility to the drug should be established. Patients receiving more than 500 mg daily should be closely observed for the signs of CNS toxicity.

PREGNANCY

Category C: Cycloserine should be given to a pregnant women only if clearly needed.

Nursing mothers: Because of the potential for serious adverse reactions in nursing infants from cycloserine, a decision should be made to whether to discontinue nursing or to discontinue the drug, taking into accounts the importance of the drug to the mother.

Usage in pediatric patients: Safety and effectiveness has not been established

PRESENTATION

Strip pack of 10 capsules. Such 6 strips are packed in a carton.

STORAGE CONDITION

Preserve in tight container below 25°C.
KEEP OUT OF REACH OF CHILDREN.

MACLEODS



NAME & ADDRESS OF MANUFACTURER:

Macleods Pharmaceuticals Ltd

Phase II, Plot No. 25 – 27, Survey No. 366, Premier Industrial Estate, .
Kachigam, Daman – 396210, India.

Off.: Atlanta Arcade, Marol Church Road, Andheri (E, Mumbai - 400 059. .

Product Registration Holder :

Zulat Pharmacy SDN. BHD.

No. 23 & 23 A, Jalan Bandar 3, Taman Melawati ,53100 , Kuala Lumpur , Malaysia.

Date of revision: Jan 2026