
**PACKAGE INSERT FOR
KORDEL'S GLUCOSAMINE
550 CAPSULE**

Name and Strength of Active

Substance(s)

Glucosamine Sulfate Potassium Chloride
750mg equivalent to 550mg Glucosamine Sulfate.

Product Description

White coloured free flowing powder in a colourless, size 00 hard gelatin capsule with:
Closed length 23.3mm +/- 0.5mm

Pharmacodynamics

Glucosamine stimulates the biosynthesis of proteoglycans, which are fundamental components of aggrecan found in articular cartilage. Aggrecan confers on cartilage its high shock-absorbing properties and resilience vital for proper joint function. Glucosamine also inhibits enzymes involved in cartilage degradation. In addition, glucosamine improves synovial fluid production and viscosity, giving better lubricant activity for the joints.

Pharmacokinetics

Following oral administration, glucosamine sulphate is well-absorbed from the gastrointestinal tract. The absolute bioavailability is 26% due to first-pass effect in the liver. Plasma levels of glucosamine reach a peak after 8-10 hours and then decline, exhibiting a half-life of 68 hours. About 10% is excreted in the urine, with about 11% found in the faeces.

Indication

As an adjuvant therapy for osteoarthritis

Recommended Dose

Adults

Take 1 capsule 3 times daily, 15 minutes before meal. Glucosamine sulphate treats the cause of osteoarthritis and pain relief can be seen only after about one week.

Route of Administration

Oral

Contraindications

Contraindicated in individuals who are hypersensitive to any of the ingredients in the product.

Warnings and Precautions

Keep out of reach of children.
Jauhkan daripada kanak-kanak.

Derived from seafood.

The effect of glucosamine on insulin sensitivity and glucose tolerance is unclear, caution is advised for diabetic patients.

Interactions with Other Medicaments

Unknown

Pregnancy and Lactation

If you are pregnant or breastfeeding, please consult your doctor or health care professional before taking this product.

Side Effects

• Cardiovascular

Peripheral oedema, tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

• Central nervous system

Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

• Gastrointestinal

Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine.

• Skin

Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

Symptoms and Treatment of Overdose

Unknown

Effects on Ability to drive and Use Machine

Unknown

Storage Conditions

Store in a dry place below 30°C.

Shelf Life

Please see label

Pack Size Available

30s, 60s, 90s

Not all pack sizes are marketed.

MAL Number

MAL06091431XC

Manufacturer

Vitaco Health (NZ) Ltd.

Cnr Kordel Place & Accent Drive,
East Tamaki, Auckland New Zealand

Product Registration Holder

Cambert (M) Sdn Bhd
8th Floor, Wisma Consplant 1,
No. 2, Jalan SS 16/4,
47500 Subang Jaya,
Selangor, Malaysia.

Date of Revision

01/06/2022