

Proj. Title: Thomson OsteoPro Insert Malaysia	
Start Date: 7/12/21	Latest Date: 4/1/24
Proj. Code: OBI-8	Size: 6.8cm(W) x 9.8cm(H)
Open Size : 13.6cm(W) x 9.8cm(H) Front & Back printing	

Applicable for all pack sizes

THOMSON OSTEOPRO	
生皮肤副作用，如红斑及瘙痒症。•孕妇及哺乳期妇女怀孕或哺乳期间服用本产品的安全性未获得证实。如果您怀孕或正在哺乳，请在服用本产品之前咨询您的医生或保健专业人士。•其他药物影响未见报道。•过量服用 - 症状与治疗未见报道。•贮存方式不要存放超过30度。避免热气和湿气。•有效期三年。 • DISTRIBUTED BY/ DIEDARKAN OLEH/ 分销商 WEST MALAYSIA/ BARAT MALAYSIA/ 西马: Zuelig Pharma Sdn. Bhd. (4013-D) No 15, Persiaran Pasak Bumi, Seksyen U8, Perindustrian Bukit Jelutong, 40150 Shah Alam, Selangor. SABAH: Apex Pharmacy Marketing Sdn. Bhd. Lot 24, Karamunsing Warehouse, Jalan Sembulan Lama, Karamunsing, 88000 Kota Kinabalu, Sabah SARAWAK: Apex Pharmacy Marketing Sdn. Bhd. AH 106-107, Batu Kawan New Township, Jalan Batu Kawa, 93250 Kuching, Sarawak. • PRODUCT REGISTRATION HOLDER AND MANUFACTURE/ PEMEGANG PENDAFTARAN PRODUK DAN PENGILANG/ 产证注册持有人和制造商 Herbal Revival Sdn. Bhd. 198701009341 (159007-X) No. 5 & 7, Jalan T5&8, Taman Industri Sungai Buloh, 47000 Sungai Buloh, Selangor Darul Ehsan, Malaysia. Tel: 603-6157 4818 Fax: 603-6157 4766  Revision Date/ Tarikh Semula: 增版日期: 4 January 2024 OBI-8	•NAME OF PRODUCT Thomson OsteoPro. •PRODUCT DESCRIPTION White body and cap, size 'O' vegetable capsule containing white to off-white powder. •INDICATION Adjunctive Therapy For Osteoarthritis. •COMPOSITION Each vegetable capsule contains 400mg Glucosamine sulfate dipotassium chloride equivalent to 300mg Fully Reacted Glucosamine sulfate of Non-Shellfish origin. Fully Reacted Glucosamine sulfate exists as a single chemical compound. In contrast, commonly available Glucosamine sulfate is a physical mixture of Glucosamine hydrochloride and potassium sulfate. Therefore, Fully Reacted Glucosamine sulfate is more uniform in dosage and is less likely to cause gastric irritation. •SOURCE OF VEGETABLE CAPSULE Hypromellose. •DESCRIPTION OF PRODUCT AND PACKAGE Pack size of 30, 60, 90, and 120 capsules in white HDPE plastic bottles. •RECOMMENDED ADULT DOSE 2 capsules, to be taken twice daily after meals. •ROUTE OF ADMINISTRATION: Oral •PHARMACODYNAMICS/ PHARMACOKINETICS Glucosamine sulfate is a natural substance that stimulates the manufacture of cartilage protein component (glycosaminoglycan) to help in joint structure repair. The anti-inflammatory effect of Glucosamine sulfate is not due to the interference with prostaglandin synthesis. Instead, Glucosamine sulfate inhibits production of superoxide radicals by macrophages and lysosomal enzymes by the liver. This absence of prostaglandin inhibition may explain its lack of ulcerogenic effect on the stomach. Following an oral administration, Glucosamine sulfate is absorbed within 15 minutes from the gastrointestinal tract and achieves peak red blood cell and plasma concentration in 4 hours. It is selectively taken up by the liver, bone and cartilage. At least 30% of the dose is concentrated in the skeletal tissue. Glucosamine is metabolised in the liver and has no active metabolites. More than 80% of the drug is recovered in the carbon dioxide of expired air and 6% and 5% are excreted in the urine and faeces respectively. •PRECAUTIONS

THOMSON OSTEOPRO	
a) Individuals who are allergic to sulfate products. b) Slow onset of action. In case of intense pain, analgesic may need to be taken. c) Keep out of reach of children. •CONTRAINDICATIONS Individuals who are allergic to glucosamine and sulfate products. •SIDE EFFECTS Cardiovascular: Peripheral oedema, tachycardia were reported in a few patients following longer clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established. Central Nervous System: Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%). Gastrointestinal: Nausea, vomiting, diarrhea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine. Skin: Skin reaction such as erythema and pruritus have been reported with therapeutic administration of glucosamine. •PREGNANCY AND LACTATION Safety of use in pregnancy and lactation has not been established. If you are pregnant or breastfeeding, please consult your doctor or health care professional before taking this product. •DRUG INTERACTION None known •OVERDOSSAGE - SYMPTOMS & TREATMENT None known. •STORAGE CONDITION Do not store above 30°C. Keep away from heat and moisture. •SHELF LIFE 3 years from date of manufacture. •产品名称 汤申护节灵 •产品说明 'O'号全白植物性胶囊内含白色或类白色粉末。•主治关节炎之辅助治疗。•成份 每粒植物性胶囊含400毫克葡萄糖胺硫酸盐二钾盐，其中含相等於300毫克，非贝壳类的全活化葡萄糖胺硫酸盐剂量。全活化葡萄糖胺硫酸盐是以单体化合物呈现。市面常见的葡萄糖胺硫酸盐则是由葡萄糖胺硫酸盐和硫酸钾混合而成。因此，全活化葡萄糖胺硫酸盐剂量，有更均匀的药性分佈，也比较不会伤	胃。•植物性胶囊源自羟丙基甲基纤维素。产品及包装说明 白色HDPE塑胶罐，内装30, 60, 90, 和120粒胶囊。•成人用量 每天二次，每次二粒胶囊餐后服食。•用途: 口服 •药理及其体内运作过程葡萄糖胺硫酸盐是天然物质，促进体内软骨蛋白的裂成，以帮助修复关节。葡萄糖胺硫酸盐的消炎效果并非因其对前列腺素合成的干扰,而是因它促使钼螯细胞和肝内溶酶酶抑制超氧自由基的产生所致。这种不抑制前列腺素的情况可说明了服食者少有胃部溃疡的现象。服用葡萄糖胺硫酸盐15分钟内，便可被消化系统吸收，并使其在红血球和血浆的浓度在四小时内达最高点，分别由肝、骨髓及软骨选取吸收。至少有30%的剂量集中在骨酸组织。葡萄糖胺在肝内新陈代谢却不产生有机的新陈代谢物。超过80%的药物可由呼出的二氧化碳中排出，而6%及5%则分别随尿液和粪便排出体外。 •警戒 (一) 对硫酸盐类产品过敏者。(二) 药力作用较缓，若疼痛持续，或可加服止痛药。(三) 放置在儿童取不到之处。•禁忌/相型作用对葡萄糖胺及对含硫酸盐产品过敏者。•副作用 心血管: 极少数患者曾在大规模的骨关节炎临床试验测试口服后被发现水肿及心跳过速。中枢神经系统: 少过1%服食者在治疗期间出现头晕、头痛及失眠现象。肠胃: 口服葡萄糖胺治疗期间极少机率会出现恶心、呕吐、腹泻、消化不良或腰部疼痛、便秘、胃灼热及厌食。皮肤: 极少数的服食者会产生