

Enzdase E.C. Tablets 5mg

Composition:

Each E.C. Tablet Contains:

Serratiopeptidase..... 5mg

Pharmacodynamics:

Serratiopeptidase is a proteolytic enzyme with antiinflammatory and antioedemic activity, it has strong enzymatic activity and excellent decomposition activity for Bradykinin. Serratiopeptidase can effectively relieve the complicated symptoms caused by inflammation, improve tissue recovery, inhibit inflammation and it can dissolve the necrotic tissue in the inflammatory lesion.

1.Relieving edema, hematoma and tumidity caused by inflammation.

It can improve the body fluid circulation of the inflammatory site complicated by surgery, trauma and inflammatory disease, and breaking down the abnormal exudate or denatured protein in the inflammatory lesion, it can relieve edema, hematoma and tumidity complicated by inflammation.

2.Dissolving sputum, pus and enhancing discharge.

Breaking down the viscous excretions remained in trachea or abdominal cavity due to bronchitis, bronchial asthma or after anesthesia, and enhancing discharge.

3.Clearing the inflammatory lesion and improving tissue recovery.

Dissolve and clear the pus cellulose and necrotic tissue in inflammatory lesion due to surgery, trauma, burn etc., and improve the regeneration of granular tissue.

Increasing the effect of antibiotics.

4.It can improve the penetration of antibiotics into infected lesion and increase the effect of antibiotics.

Pharmacokinetics :

Serratiopeptidase when consumed in unprotected form is destroyed by acid in the stomach.

However, enteric coated granules, enable the enzyme to pass through the stomach unchanged,

and be absorbed in the intestine. It is found negligibly in urine suggesting that it is transported directly from the intestine into the blood stream.

Indications:

Inflammation after surgery or trauma, inflammation of nasal cavity, cystitis, epididymitis, peripheral inflammation of teeth , alveolar abscess; Inadequate expectoration of sputum in bronchitis, bronchial asthma or after anaesthesia.

Adverse Effects:

1. Exanthema, rubefaction are rare and usually disappear rapidly after discontinuation of Serratiopeptidase.

2. Diarrhea, loss of appetite, gastric disturbance, nausea, vomiting rarely occur.

3. Rarely,bleeding tendency eg. epistaxis and bloody sputum may occur.

Precautions/Warnings:

1. It should be administered with caution in patients with blood coagulation abnormalities, severe hepatic and renal insufficiency.

2. It should be administered concomitantly with antibiotics (or chemotherapy) in patients complicated with various bacterial infections.

Use in Pregnancy and Lactation:

No well-controlled studies are available regarding the use of Serratiopeptidase in pregnancy and lactation.

Contraindications:

It is contraindicated in patients who are hypersensitive to it.

Dosage :

Oral administration.

15 ~ 30 mg daily; 1 ~ 2 tablets to be taken 3 times daily after food.

As it is an enteric film coated tablet, please do not chew but to be swallowed whole.

Symptoms and treatment of overdosage:

No data available on overdosage with Serratiopeptidase. In the event of an overdosage, the stomach should be emptied and symptomatic treatment should be given.

Drug interactions:

Since the concomitant use of Serratiopeptidase with an anticoagulant may intensify the anticoagulant effect, Serratiopeptidase should be administered with caution and under close monitoring.

Storage:

Store below 30°C. Protect from direct light.

Shelf-Life: 3 years

Pack sizes:

Blister pack: A box of 10 strips of 10 tablets per strip.

Description:

A light peach, round shaped enteric coated tablet.

The uncoated tablet size is 9/23" D.C.

The film coated tablet weight is 132mg

Appearance is a light peach enteric coated tablet.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM
YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Revised Date: 16/08/2011

Product Registration Holder:

Kireen Pharmaceutical Sdn. Bhd.

No. 9 , Jalan Kempas 4, Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.

Manufactured by:



C.C.P.C.

CHINA CHEMICAL & PHARMACEUTICAL CO., LTD.

Hsinfong Plant: 182-1 Keng Tze Kou, Hsinfong, Hsinchu, Taiwan, R.O.C. (A)

98.07.01製

101.05.22修改內容

101.05.25修改內容

101.05.28輸出生產

114.10.01修改內容

130x310mm