

Cartril-S™

Glucosamine HCl

NAME OF THE MEDICINAL PRODUCT

Cartril-S.

Gelatin for capsule: Bovine source certified with Halal certification.

COMPOSITION

Cartril-S 1500mg Powder

Each sachet contains 1417.5 mg of Glucosamine HCl equivalent to Glucosamine content in 1500 mg Glucosamine Sulphate.

Cartril-S 500mg Capsule

Each capsule contains 472.5 mg of Glucosamine HCl equivalent to Glucosamine content in 500 mg Glucosamine Sulphate.

Preservatives: None

DESCRIPTION

Cartril-S 1500mg Powder: Free flowing white to off white colour odourless and tasteless powder.

Cartril-S 500mg Capsule: Size1, hard gelatin capsule with red cap and red body containing white to off white powder. Printed 'Cartril S' and '500'.

PHARMACODYNAMIC

Glucosamine possesses arthrophic metabolic activity by stimulating the anabolic metabolism of osteocartilaginous tissues, through stimulation of biosynthesis of the mucopolysaccharides (which are essential components of the cartilage ground-substances) and the bone mesenchymal tissues and improvement of the calcium intake. Lubricant activity through improvement of synovial fluid by increasing synovial fluid production.

PHARMACOKINETICS

Absorption

Glucosamine sulphate is well absorbed (with a proportion close to 90%) following oral administration.

Protein binding

It is known that glucosamine and its metabolites bind to α - and β -globulins.

Distribution

Mainly distributed to bone tissues and articular cartilages.

INDICATIONS

As adjunct therapy in osteoarthritis.

CONTRAINDICATIONS

Hypersensitive to Glucosamine. The powder for oral solution contains aspartame and therefore contraindicated in patients with phenylketouria. Contraindicated in patients allergic to shellfish.

DRUG INTERACTION

Drug interaction with Oral Hypoglycaemic Agent, may reduce the oral hypoglycaemic agent effectiveness. For examples; metformin, tolbutamide, rosiglitazone and etc.

Glucosamine is likely safe in patients with well controlled diabetes (HbA1c less than 6.5%) taking one or two oral diabetics medications or controlled by diet only. In patients with higher hbA1c levels or those taking insulin, monitor blood glucose levels closely.

PRECAUTIONS/WARNINGS

The administration in patients with severe hepatic or renal insufficiency should be made under medical supervision. Glucosamine is not meant for replace NSAIDs in cases of intense pain, as this is a causal therapy. It is advisable to take an anti-inflammatory drug in the initial treatment of osteoarthritis together with glucosamine.

No effects on the ability to drive or to operate machines is expected.

Pregnancy and Lactation

The animal studies did not show any evidence of unfavourable effects on the reproductive functions or on lactation. In the absence of such studies in humans, the product during pregnancy and

lactation should be used only under medical control. Administration during the first three months of pregnancy must be avoided.

DOSAGE

Cartril-S 1500mg Powder:

The content of one sachet (dissolved in a glass of water) should be taken once daily, preferably at meals.

Duration of treatment: 3 months.

(The treatment course may be repeated at intervals of 2 months).

Cartril-S 500mg Capsule:

Adult: One capsule taken twice daily for at least 6 weeks (or according to medical prescription).

Duration of treatment: 3 months.

(The treatment course may be repeated at intervals of 2 months).

OVERDOSE

No cases of accidental or intentional overdose are known. The animal acute and chronic toxicological studies indicate that toxic effects and symptoms toxicity are not likely to occur, even after high overdoses.

UNDESIRABLE EFFECTS

Undesirable effects have been observed in low proportion of patients. They were transient, of minor entity, and included gastric discomfort and pain, meteorism, constipation, diarrhoea. Effects of hypersensitivity have been reported in some patients and included cutaneous rashes with pruritus and erythema.

The patient should inform his doctor of any other undesirable effect during the treatment.

Cardiovascular

Peripheral oedema, tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

Central nervous system

Drowsiness headache, insomnia have been observed rarely during therapy (less than 1%).

Gastrointestinal

Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been describe rarely during oral therapy with glucosamine.

Skin

Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

PACKING/PACK SIZE

Cartril-S 1500mg Powder: 3.95gm x 30 sachets

Cartril-S 500mg Capsule: 10 x 10's (Blister) & 60's (Bottle)

STORAGE CONDITIONS, USER INSTRUCTIONS, AND PHARMACEUTICAL PRECAUTIONS

Store in a dry place below 30°C.

Protect from moisture and heat.

Keep out of reach of children.

Jauhi daripada kanak-kanak.

Derived from Seafood

Shelf-life: Please refer to outer package

Do not use the product after the expiry date indicated on the label.

Avoid exposure to excessive heat.

The expiry date applies to the correctly stored product in the intact package.

Product Registration Holder:

DUOPHARMA MARKETING SDN BHD.

Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.

Manufacturer:

DUOPHARMA MANUFACTURING (BANGI) SDN BHD.

Lot No. 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.



Revision date: April 2019

1500008537 N.1

150mm(w) x 211mm (h)



Black 100%