

Seven Seas Jointcare Glucosamine 500mg Capsule (Package Insert)

SEVEN SEAS JOINTCARE GLUCOSAMINE 500MG CAPSULE

DESCRIPTION

Reddish orange opaque and white opaque capsule with SEVEN SEAS logo printed on the cap and "GLUCO 500" printed on the body.

COMPOSITION

Each capsule contains Glucosamine Potassium Sulfate 663.20 mg (equivalent to 500mg of Glucosamine Sulfate).

List of excipients

- *Corn Starch*
- *Microcrystalline Cellulose*
- *Talc*
- *Sodium starch Glycolate*
- *Magnesium stearate*

PHARMACODYNAMICS

Glucosamine possesses antiarthrosic-arthrotrophic activities. It stimulates the anabolic metabolism of osteo-cartilaginous tissues via stimulation of the biosynthesis of the mucopolysaccharides (which are the essential components of the cartilage ground substances) and of the bone mesenchymal tissues. Glucosamine also acts to improve the viscosity of synovial fluid by increasing synovial fluid production, thereby providing lubricant activity.

PHARMACOKINETICS

Absorption

Glucosamine sulfate is well absorbed (with a proportion close to 90%) following oral administration.

Distribution

Mainly distributed to bone tissues and articular cartilages.

INDICATIONS

Adjunctive therapy in osteoarthritis.

CONTRAINDICATIONS

Contraindicated in patients sensitive to glucosamine sulfate and shellfish.

WARNINGS AND PRECAUTIONS

- Keep out of reach of children.
- Seven Seas Jointcare Glucosamine 500mg Capsule is not meant to replace NSAIDs in cases of intense pain, as this is a causal therapy. It is advisable to take an anti-inflammatory drug in the initial treatment of osteoarthritis together with glucosamine sulfate.
- Glucosamine may increase insulin resistance. Those with type 2 diabetes and those who are overweight and have problems with glucose tolerance should have their blood sugars carefully monitored if they consume glucosamine.
- Because of insufficient data, children, pregnant women and lactating mothers should avoid using glucosamine.
- **Derived from SEAFOOD (SHELLFISH)**
- **Use in pregnancy and lactation**
To be taken under medical supervision.
- **Paediatrics**
Not recommended.
- **Geriatrics**
To be taken under medical supervision.
- **Drug interactions**
None known
- **Incompatibilities**
None known

ADVERSE EFFECTS/ UNDESIRABLE EFFECTS:

Cardiovascular

Peripheral oedema, tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

Central nervous system

Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

Gastrointestinal

Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine.

Skin

Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

Immune system

Allergic reactions including rash, itching, hives, swelling of face, lips tongue or throat.

DRUG INTERACTIONS

Glucosamine may increase insulin resistance and consequently affect glucose tolerance. Diabetics who are under medical advisement, decide to use glucosamine will need to monitor their blood glucose and may need to adjust the doses of the medications they take to control blood glucose. This needs to be done under medical supervision. Interactions between glucosamine and warfarin may result in the magnification of the anticoagulant effects of warfarin. If you are taking warfarin, talk to your physician before using this product.

SYMPTOMS OF OVERDOSE AND TREATMENT

None known.

DOSAGE AND ADMINISTRATION

Adults : For mild to moderate osteoarthritis symptoms: Oral; 1 capsule 3 times daily for 6 weeks

For severe osteoarthritis symptoms: Oral; 1 capsule 3 times daily for 8 weeks.

Maintenance therapy should be followed for 3-4 months by administration of 1 capsule 2 times daily. Do not exceed the stated recommended daily dose.

Storage: Store below 30°C. Protect from light and moisture.

Presentation/Packing: Glass bottles of 30s, 60s, and 90s.

Product License Holder :

Allpac Solutions Sdn Bhd (457587-X)

121, Jalan Tunku Abdul Rahman

(Jalan Kuala Kangsar),

30010 Ipoh, Perak, Malaysia.

Distributed by:

Zuellig Pharma (M) Sdn Bhd (4013-D)

Manufactured for Seven Seas by:

HOVID Bhd. (58476-A)

Lot 56442, 7 1/2 Miles, Jalan Ipoh/Chemor,

31200 Chemor, Perak, Malaysia

Date of Revision: September 2024