



止跌
痛打
如风
神湿

成份/Composition:

每毫升含有/Each ml contains:

冬青油	Methyl Salicylate	60%w/v
薄荷	Menthol	10%w/v
樟脑	Camphor	3%w/v

外观: 带有特征气味的棕色油膏。

主治: 缓解由风湿引起的肌肉和关节疼痛。

用法: 将此油搽于患处, 每天3至4次。

药效率: 请参阅英文版本。

禁忌: 对此产品中任何成份过敏者。

警戒和注意事项:

禁止于2岁以下的婴儿。用于较年长儿童时需谨慎。将含有樟脑的产品涂于儿童的鼻孔是很危险的。此法少量应用也可能导致休克。此产品含有水杨酸甲酯, 当涂抹或擦在皮肤时, 可以通过皮肤渗透进入血液。对于服用华法林的患者, 当因肌肉或关节疼痛时过度涂抹于皮肤时可能会增加出血的可能性。避免与眼睛和粘膜接触。不要涂抹于伤口或受损皮肤。避免在怀孕和哺乳期间过度使用。不要大面积涂抹在身体或用绷带包扎或使用加热垫。在涂抹于较大区域的皮肤前, 涂抹少量在有限的皮肤区域以观察是否有不适或痒痒。若刺激反应持续, 请停止使用。

外用忌食。

侧面效应/不良反应: 皮肤刺激, 发红或肿胀, 灼热或刺痛感。不良反应很少见。可能会出现头晕和恶心的症状。

对驱动和使用机器的能力的影响: 未知。

妊娠和哺乳期间使用声明: 请参阅英文版本。

贮藏: 储存于30°C以下。放在儿童不能接触的地方。

Product Description: Brown-coloured oil with characteristic odour.

Indication: To relieve muscle and joint pain related to rheumatic pain.

Direction for use: Rub on site of pain 3 to 4 times daily.

Pharmacodynamics:

Camphor is a stimulant, rubefacient, antipruritic and feebly antiseptic. When in combination with other ingredient like menthol and methyl salicylate, it becomes ideal for neuralgia and other painful areas.

Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Methyl Salicylate is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins, and cause platelet dysfunction. Methyl salicylate is used topically as a counter-irritant. Upon application, it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscle or joints.

Pharmacokinetics:

Camphor is readily absorbed from all administration sites. Camphor is well absorbed after dermal exposure. It is hydroxylated in the liver to yield hydroxycamphor metabolites which are then conjugated with glucuronic acid and excreted in the urine. Camphor crosses the placenta.

After absorption, menthol is excreted in the urine and bile as a glucuronide.

Methyl Salicylate may be absorbed through intact skin. The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity. Absorption characteristics of salicylates vary with the dose, formulation and route of administration. Percutaneous absorption is enhanced by exercise, heat, occlusion or disruption of the integrity of the skin or application to large areas of skin. Both the rate and extent of absorption increases after repeated application; increasing the bioavailability. Methyl salicylate is extensively metabolized to salicylic acid in the dermal and subcutaneous tissues after topical application. At therapeutic levels, the half-life of salicylates is 2-4 hours. As salicylate level increase into the toxic range, the half-life can be greater than 18 hours.

Contraindications: Patient with known hypersensitivity to any of the ingredients in the product.

Warnings and Precautions:

This product is contraindicated in infants below 2 years of age. Caution must be exercised when older children are treated. It is dangerous to place any camphor containing product into the mouth of children. A small amount applied there may cause irritation or collapse.

This product contains methyl salicylate and when applied or rubbed on the skin can be absorbed through the skin into the blood.

For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

Avoid contact with the eyes and mucous membrane. Do not apply to open wounds or damaged skin. Avoid excessive use during pregnancy and lactation. Do not apply over large areas of the body or bandage or use with heating pads.

Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to a larger skin area. Discontinue use if irritation develops.

For external use only. Not to be taken.

Effects on Ability to Drive and Use Machines: Not known.

Interactions with Other Medicaments:

This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joints pains may increase the chances of bleeding.

Statement on Usage During Pregnancy and Lactation:

Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child.

SIDE EFFECT / ADVERSE REACTION: Skin irritation, redness or swelling, burning or stinging sensation. Adverse reaction is rare. Symptoms of dizziness and nausea may occur.

Symptoms and Treatment of Overdose: Mild chronic salicylate intoxication or salicylism usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache and confusion and may be controlled by reducing the dosage.

Storage Conditions: Store below 30°C. Keep out of reach of children. Jauhkan daripada kanak-kanak.

Packing Size: 24ml, 23ml, 57ml and 60ml

Shelf Life: 3 years from date of manufacture.

1/AL07031279XC

给药途径: 外用

药动学: 请参阅英文版本。

与其他药物的相互作用: 请参阅英文版本。

过量症状和治疗: 请参阅英文版本。

保质期: 自生产日期起3年。

Route of Administration: Topical

TIGER HILL MEDICATED OIL

MINYAK UBAT GUNUNG HARIMAU

Manufactured by:
WELLMETECH PHARMACEUTICAL INDUSTRIES SDN BHD
Lot 8058, Kota Samarahan Industrial Park, 94300 Kota Samarahan, Sarawak, Malaysia.

Marketing Authorization Holder:
KIAN FENG TRADING
81, Laluan Rokam 13, Pekan Razaki, 31350 Ipoh, Perak

Distributed by:
EYH FONG TRADING
10, Jalan Rapat Baru 21, Kawasan Perindustrian Ringan Kinta Jaya, 31350 Ipoh, Perak