

FUCID OINTMENT 2% W/W

COMPOSITION:

Each gram of ointment contains: Sodium Fusidate (Micronised) 20mg or 2.0% w/w

DESCRIPTION:

White to pale yellow, smooth ointment.

PHARMACODYNAMIC:

Sodium fusidate is an antibiotic derived from *Fusidium coccineum* and exerts powerful antibacterial activity against a number of gram-positive organisms. Staphylococci, including the strains resistant to penicillin or to other antibiotics, are particularly susceptible to sodium fusidate.

PHARMACOKINETIC:

In vitro studies show that fusidic acid can penetrate intact human skin. The degree of penetration depends on factors such as the duration of exposure to fusidic acid and the condition of the skin. Fusidic acid is excreted mainly in the bile with little excreted in the urine.

INDICATION:

Sodium fusidate ointment is indicated for the treatment of skin infections caused by staphylococci, streptococci, *Coryne-bacterium minutissimum* and other organisms sensitive to sodium fusidate. The most important indications being:- Impetigo, Boils, Carbuncles, Paronychia, Infected wounds, Sycosis barbae, Hidradenitis, Erythrasma, Folliculitis.

RECOMMENDED DOSAGE:

Apply to the affected area 2-3 times a day, generally for a period of 7 days. It can be used with or without a covering dressing.

CONTRAINDICATION:

Hypersensitivity to the preparation.

Infection caused by non-susceptible organisms, in particular, *Pseudomonas aeruginosa*.

WARNING AND PRECAUTION:

Sodium fusidate has been shown to cause conjunctival irritation. The ointment should not be used in or near the eyes. Bacterial resistance has been reported to occur with the use of sodium fusidate applied topically. As with all topical antibiotics, extended or recurrent application may increase the risk of contact sensitisation and the development of antibiotic resistance.

DRUG INTERACTION:

None known.

PREGNANCY AND LACTATION:

There is evidence to suggest that when given systemically, fusidic acid can penetrate the placental barrier. The use of Fucid Ointment in pregnancy requires that the potential benefits be weighed against the possible hazards to the foetus.

Safety in nursing mothers has not been established. When sodium fusidate has been given systemically, levels have been detected in breast milk but with topical use the possible amount of drug present is unlikely to affect the infant.

SIDE EFFECTS:

Hypersensitivity reactions to the active ingredient in the form of skin rashes, mild stinging and irritation on application have been reported rarely.

OVERDOSAGE:

Not applicable

PRESENTATION:

Aluminium tubes of 5 gm, 10 gm and 15 gm

STORAGE:

Store below 30°C.

For external use only.

Keep containers tightly closed. Keep out of reach of children. Jauhi dari kanak kanak.

MANUFACTURED BY: NORIPHARMA SDN. BHD. 200701034604 (792633-A)

Lot 5030, Jalan Teratai,

5 ½ Mile Off Jalan Meru,

41050 Klang, Selangor, Malaysia.

DATE OF REVISION: December 2022