

133 mm.



FluiFort®

Syrup

International Non-proprietary Name

Carbocysteine Lysine Salt Monohydrate.

Composition

100 ml of **FluiFort**® Syrup contains Carbocysteine Lysine Salt Monohydrate, (Carbocysteine Lysine Salt 9 g) equivalent to Carbocysteine 5 g.

Description

A clear, brilliant, yellowish brown liquid.

Therapeutic Properties

FluiFort (Carbocysteine Lysine Salt Monohydrate) is a muco-regulator which exerts a specific *metabolic action* on the serous and mucous cells of the respiratory epithelium. It has no effect on the mucus itself.

The main demonstrated effect of **FluiFort**® is the increase of sialomucin contents in the mucus, restoring the proper sialo- to fuco-mucin proportion. This directly results in restored physiological mucus viscosity, elasticity and adhesion, and indirectly in restoring mucus hydration and mucociliary clearance.

The salification of Carbocysteine with Lysine results in greater solubility and biological availability, which allows to decrease the number of daily administrations while maintaining the wanted effects, and with an increase of the post-dosing effect typical of metabolic agents. Once the metabolic mechanism is activated – that is, after one or a few days of administration, depending on the specific condition of the patient – it remains in effect even in the absence of drug administration (post-dosing carry-over effect).

Pharmacokinetics

Carbocysteine Lysine Salt Monohydrate has a greater water solubility and hence improved bio-availability. Therefore it is possible to administer **FluiFort**® once daily in adults and twice daily in children.

The active ingredient is rapidly and completely absorbed after oral administration (plasma half-life 1.5 hour). It is mainly excreted via the urinary route; 30-60% is excreted unmodified; the remainder in the form of various metabolites. Bio-availability studies have shown that up to 17.5% of the administered dose can be found in the bronchial secretions.

Indications

FluiFort® is indicated for the treatment of acute and chronic respiratory tract disease accompanied by excessive mucous.

Recommended Dose

Adults:

30 ml once daily or 15 ml twice daily.

Children:

- Not recommended in neonates and infants.
- **2 to 5 years of age:** 2.5 ml twice daily.
- **over 5 years of age:** 5 ml twice daily.

Administration in children should be done under careful medical monitoring.

The dosage may be increased to administration three times daily (t.i.d.) in both children and adults according to medical prescription.

Contraindications

History of ascertained hypersensitivity to the product; gastric or duodenal ulcer.

Contraindicated in children below 2 years of age.

Precautions

When treating diabetic patients or in case of low-calorie diets it should be considered that **FluiFort**® Syrup contains sucrose.

Use During Pregnancy and Lactation

Although the product is neither teratogenic nor mutagenic, and did not appear to have negative effects on the reproductive function in animal studies, it is not recommended for use during the first trimester of pregnancy.

As it is not known whether carbocysteine is excreted in milk, its use during lactation is not advised.

Overdosage

In case of overdosage with enhancement of side effects, gastric lavage may be useful, followed by additional treatment under medical supervision. No specific antidote exists.

Side Effects

The clinical trials and pharmacovigilance initiatives conducted after commercialisation show that **FluiFort**® is well tolerated. Occasionally, gastrointestinal symptoms such as gastralgia, nausea, diarrhoea, mild to moderate and transient dizziness, and very rarely skin rashes have been reported. These side effects disappear upon reduction of the dosage.

Immune System Disorders:

Anaphylactic / anaphylactoid reaction

Skin and Subcutaneous Tissue Disorders:

Severe cutaneous adverse reactions (SCAR) e.g. erythema multiforme, Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN). In most of these cases reported at least one other drug was administered at the same time, which may have possibly enhanced the described mucocutaneous effects.

Drug Interactions

No incompatibilities are known between **FluiFort**® and the most commonly used medications for the treatment of disorders of the upper and lower airways.

Packaging

Bottles containing 60 ml and 120 ml.

Storage Conditions

Store in a cool, dry place below 30 °C.

Keep out of reach of children / Jauhi dari kanak-kanak

Manufactured under licence for:

Eurodrug Laboratories S.A. (Belgium)

by M & H Manufacturing Co., Ltd.,

Samutprakarn, Thailand.

Original Research Product of Dompé Farmaceutici S.p.A. (Italy).

Product Registration Holder:

Eurodrug Laboratories (M) Sdn Bhd

T2-21-10, The Square, One city,

Jalan USJ 25/1C, 47650 Subang Jaya,

Selangor, Malaysia

Date of revision text:

June 2019



02MY27199.A01
20180822AH
N0901

183 mm

Black