

Acti// -Rub Liniment

Active Ingredient:

Methyl Salicylate 25%w/v

Presentation:

Bottle of 30ml.

Product Descriptions:

Colourless, clear and non-greasy solution with characteristic odour.

Pharmacology:

Methyl Salicylate is used as a rubefacient in deep heating liniments and principally for the symptomatic treatment of mild to moderate pain, rheumatic and inflammatory conditions. It increases blood flow to the area, provides feeling of warmth and gives rapid relief of aches and pains.

Indications:

Activ-Rub is used for the temporary relief of pain associated with muscular cramp, bruises, sprains and strains, muscle aches, rheumatism, arthritis and other similar conditions.

Directions of Use:

Apply a small amount onto affected areas and gently massage until absorbed. Repeat 3 times a day or reapply as needed for continued relief.

Symptoms and Treatments of Overdose:

Symptoms of mild chronic Salicylate intoxication or salicylism include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache and mental confusion. Symptoms of severe intoxication include hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and metabolic acidosis. Depression of the central nervous system may lead to coma, cardiovascular collapse and respiratory failure. In children, drowsiness and metabolic acidosis commonly occur, hypoglycaemia may be severe.

In acute overdosage, the stomach should be emptied by emesis or by aspiration or lavage. Patient should be encouraged to drink plenty of fluids. In severe intoxication, forced alkaline diuresis may be required and should be continued until the plasma Salicylate concentration is less than 350 mcg/ml in adults. Then, intravenous fluid may be stopped and patient is encouraged to take fluids by mouth. Plasma electrolytes, especially potassium and acid-base balance should be monitored regularly.

Acidaemia must be corrected by infusion of Sodium Bicarbonate before starting forced diuresis. In the presence of cardiac or renal impairment or in very severe intoxication, haemodialysis or peritoneal dialysis may be necessary.

Contraindication:

Do not use if hypersensitive to Methyl Salicylate.

Precautions:

For external use only. Avoid in contact with eyes, nose and mouth. Not to be taken. If symptoms persist, please consult a doctor.

This product contains Methyl Salicylate and when applied or rub onto the skin, it can be absorbed through the skin into blood. For patients taking Warfarin, excessive application onto the skin for muscular or joint pains may increase the chances of bleeding.

Do not apply onto open wounds.

For people with sensitive skin, the applied skin may appear red but this will soon fade away.

As this preparation contains benzyl alcohol, its use should be avoided in children under 2 years of age. Not to be used in neonates.

Undesirable Effects:

Occasionally, skin sensitivity may happen in some sensitive people. Skin rash, itching and redness have also been reported.

Interactions with Other Medicaments:

Methyl Salicylate may enhance the hypoprothrombinemic effect of Warfarin and increase the risk of bleeding complications with this agent.

Storage:

Store below 30°C. Protect from light. Keep cap tightly closed after opening. Keep out of reach of children. Jauhkan daripada capaian kanak-kanak.

Shelf Life:

3 years from the date of manufacture.

Manufacturer and Product Registration Holder

WINWA MEDICAL SDN. BHD.

1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

MAL07082989X

Date of Revision: 15 July 2020

X.EAR-I.04

Size: 80mm x 100mm



Dark Blue (Pantone 2767U)