103,00 mm

OSTEOFREE 500
Capsule

Description
White powder filled in No.0 Orange (body)- Ivory (cap) capsule.

Composition
Each capsule contains:
Glucosamine Sulfate 2KCl 666.49 mg equivalent to Glucosamine Sulfate ..... 500 mg

Inactive Ingredients:
Avicel PH 102, Explotab, and Magnesium Stearate.

Pharmacology
Glucosamine possesses arthrotrophic metabolic activity by stimulating the anabolic metabolism of osteocartilaginous tissue through stimulation of the biosynthesis of the mucopolysaccharides (which are the essential components of the cartilage ground-substances) and of the bone mesenchymal tissue. It also possesses lubricant activity through improvement of synovial fluid viscosity by increasing synovial fluid production.

Pharmacokinetics
Orally administered glucosamine is rapidly absorbed, highly distributed and efficiently cleared.
The gut rather than liver is mainly responsible for the first pass metabolism since reduced bioavailability is observed after oral doses.
Distribution:
Total Protein Binding:
1. Glucosamine is not protein-bound but rather incorporates into plasma proteins (primarily globulins).
2. Unbound glucosamine did not bind to plasma proteins of humans, dogs, or rats.

Excretion:
1. Breast Milk: Breast feeding: Scientific evidence for the safe use of Glucosamine Sulfate during lactation is not available
2. Kidney: Renal excretion: 10 %
3. Feces: 11 %

Elimination Half-Life: 70 Hours

Indications
Adjunctive/adjuvant therapy in osteoarthritis.

Recommended Dose and Administration
Adults: For light to moderate osteoarthritis symptoms: Oral: 1 capsule twice daily for 6 weeks, taken 15-30 minutes before meal time.
For severe osteoarthritis symptoms: Oral: 1 capsule 3 times daily for 8 weeks, taken 15-30 minutes before meal time .
Maintenance therapy should be followed 3-4 months by administration of 1 capsule twice daily.

Contraindications
- Hypersensitivity to glucosamine.
- Contraindicated in patients allergic to shellfish.

Warnings and Precautions
- Derived from seafood.
- The effect of glucosamine on insulin sensitivity and glucose tolerance is unclear, caution is advised for diabetic patients.
- Glucosamine is not meant to replace NSAIDs in cases of intense pain as this is a causal therapy. It is advisable to take an anti-inflammatory drug in the initial treatment of osteoarthritis together with Glucosamine Sulfate.
- Glucosamine Sulfate is to be taken under medical supervision in pregnant and lactating mothers, and in geriatric patients.

Interactions with Other Medicaments

Antidiabetic Agents	: Reduced antidiabetic agent effectiveness. Glucosamine is likely safe for patients with diabetes that is well-controlled with diet only or with one or two oral antidiabetic agents (HbA1c less than 6.5%). In patients with higher HbA1c concentrations or for those requiring insulin, closely monitor blood glucose concentrations.
Doxorubicin	: Reduced doxorubicin effectiveness. Avoid concomitant use of glucosamine with doxorubicin.
Etoposide	: Reduced etoposide effectiveness. Avoid concurrent use of glucosamine and etoposide.
Teniposide	: Reduced teniposide effectiveness. Avoid concurrent use of glucosamine and teniposide.

Pregnancy and Lactation
Safety in pregnancy has not been established. Consult physician or pharmacist before administering to pregnant women or women of child-bearing potential.
Breast feeding: Scientific evidence for the safe use of Glucosamine Sulfate during lactation is not available.

Side Effects
Gastrointestinal disturbances such as nausea and vomiting are rarely reported.

Symptoms and Treatment at Overdose
None reported so far.

Storage
Store below 30°C. Protect from light.
Keep out of reach of children.

Shelf Life
36 months from the date of manufacture.

Pack Size
Box contains 10 strips of 10 capsules.

Imported and marketed by:
The Zyfas Medical Co
No 15, Jalan Mega 1/5,
Taman Perindustrian Nusa Cemerlang,
79200 Iskandar Puteri,
Johor, Malaysia

Date of Revision : Oct 2025

Manufactured by :
Landson
PT. PERTIWI AGUNG
Pharmaceutical Industry
A member of MENSA GROUP
Jl. DDN Sukadanau, Cikarang Barat
Bekasi 17843 - Indonesia

Kode : 5 CO 02 ABA

320,00 mm

3.00 mm
2.00 mm
3.00 mm
18.00 mm

Produk	: **OSTEOFREE [export]**
- negara	: Malaysia
- kemasan	: Brosur
- kode kemasan	: 5 CO 02 ABA
- ukuran	: 320 x 103 mm
- bahan	: HVS 60 g
- coating	: Biasa
- letter	: Arial Arial Narrow Switzerland Condensed Times New Roman Calligrapher
Warna	
- dasar	: putih + hitam
- judul	: hitam
- logo Landson	: hitam
- teks judul	: putih
- teks	: hitam
Perubahan	: -
Dibuat oleh	: Div. Export
Revisi	: -
- tanggal	: 29/10/25

Catatan :
- *pembagian warna ini hanya sebagai print out*