

Vinodine™

ANTISEPTIC GERMICIDAL SPRAY

For Medical Practitioner or Hospital or Laboratory use only

Vinodine Spray (Povidone-Iodine 5.0% w/w)

Composition:	% w/w
Each 100g contains:	
Povidone-Iodine USP	5.0
(0.5% w/w available Iodine)	
Inert Solvent and propellant q.s to	100.0
Excipients:	
Iso-propyl Alcohol	80.0

Product Description

Deep brown coloured liquid spray with characteristic odour of Iodine filled in pressurized Aluminium container equipped with valve.

Product Indication / Usage

Antiseptic for the treatment and prevention of infection in wounds including ulcers, burns, cuts and other minor injuries.

Dose / Use Instruction

For topical use only. Adults and children aged 2 years and over:

Shake the can well before use. Spray the required area from a distance of 15-25cm (6-10 inches) until a dusting of powder is deposited. If necessary, the treated area may be covered with a dressing.

Contra – Indication

Hypersensitivity to iodine. Vinodine Spray should not be used in serous cavities. Do not use this product regularly if the patient suffers from thyroid disorders or is receiving concurrent lithium therapy. Do not use on children under two years of age.

Warning/ Precautions

Avoid inhaling or spraying into the eyes. Avoid use on patients with renal impairment. Should evidence of local irritation or sensitivity occur, use of the product should cease. If no improvement occurs, a doctor should be consulted. Special caution is needed when regular applications to broken skin are made to patients with pre-existing renal insufficiency.

Size: W115mm x H90mm

Drug Interaction

Use with concurrent lithium therapy has been shown to exhibit additive hypothyroidic effects.

Absorption of iodine from povidone iodine through either intact or damaged skin may interfere with thyroid function test. Contamination with povidone iodine of several types of tests for the detection of occult blood in faeces or blood in urine may produce false-positive results.

Side Effects/Adverse Reaction

Iodine is absorbed through burns and broken skin and to a lesser extent through intact skin. Following prolonged application of Vinodine Spray to severe burns or large areas of denuded skin, systemic effects such as metabolic acidosis, parametrical renal impairment and thyroid dysfunction may occur.

Signs and Symptoms of Overdose and Treatment

Excess iodine can produce goitre and hypothyroidism or hyperthyroidism. Systemic

absorption of iodine after repeated application of povidone iodine to large areas of wounds or burns may lead to a number of adverse effects: metallic taste in mouth, increased salivation, burning or pain in the throat or mouth, irritation and swelling of the eyes, pulmonary oedema, skin reactions, gastrointestinal upset and diarrhoea, metabolic acidosis, hypernatraemia and renal impairment.

Treatment: In the case of deliberate or accidental ingestion of large quantities of Vinodine, symptomatic and supportive treatment should be provided with special attention to electrolyte balance and renal and thyroid function.

Storage Condition

Store below 30°C and protect from light.

Shelf Life

36 Months

Presentation:

Aluminium Container containing

75 g and 100g of 5.0 % Povidone-Iodine solution & is fitted with valve & with actuator.

Manufactured by:

**M/s. Midas Care
Pharmaceuticals Pvt. Ltd.**
B-16, MIDC, Waluj, Chhatrapati Sambhajanagar - 431136, India.
Mfg Lic No. : AD/009

Product Registration

Holder/Imported by:

UNIMED SDN BHD
53, Jalan Tembaga SD 5/2B
Bandar Sri Damansara 52200,
Kuala Lumpur.
Revised Date: 01/12/2025