

1 Tradename

PATADAY Ophthalmic Solution 0.2%w/v

2 Description and composition

Pharmaceutical form(s)

Eye drops, solution

Colorless to light yellow solution

Active substance(s)

2.22 mg of olopatadine hydrochloride in one mL solution (0.2%)

Excipients

Benzalkonium chloride, edetate disodium, povidone, sodium chloride, dibasic sodium phosphate (anhydrous), sodium hydroxide and/or hydrochloric acid, purified water.

3 Indication

PATADAY is indicated for the temporary prevention of itching of the eye due to allergic conjunctivitis.

4 Dosage regimen and administration

Dosage regimen

Adults

The recommended dose is one drop in each affected eye once a day.

Pediatric Use

Safety and effectiveness in pediatric patients below the age of 3 years have not been established.

Geriatric Use

No overall differences in safety and effectiveness have been observed between elderly and younger patients.

Method of administration

- For topical ocular use only. Not for injection or oral use.
- To avoid contamination, the dropper tip should not touch any surface. The dropper tip should also not come into contact with the eye as this may cause injury to the eye.
- When using nasolacrimal occlusion or closing the eyelids for 2 minutes, the systemic absorption is reduced. This may result in a decrease in systemic side effects and an increase in local activity.
- The bottle should be kept tightly closed when not in use.
- If more than one topical ophthalmic medicinal product is being used, the medicinal products must be administered at least 5 minutes apart. Eye ointments should be administered last.
- Patients should be advised not to wear a contact lens if their eye is red.

- Pataday should not be used to treat contact lens related irritation.

5 Contraindications

Hypersensitivity to the active substance or to any of the excipients.

6 Warnings and precautions

Special excipients

Pataday contains benzalkonium chloride which may cause eye irritation and may possibly discolor soft contact lenses. Contact lenses must be removed before administration of Pataday and reinserted at least 15 minutes later.

7 Adverse drug reactions

Tabulated summary of adverse drug reactions from clinical trials

Adverse drug reactions from clinical trials (Table 7-1) are listed by MedDRA system organ class. Within each system organ class, the adverse drug reactions are ranked by frequency, with the most frequent reactions first. Within each frequency grouping, adverse drug reactions are presented in order of decreasing seriousness. In addition, the corresponding frequency category for each adverse drug reaction is based on the following convention (CIOMS III): very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$).

Table 0-1 Adverse drug reactions in clinical trials

System Classification	Organ	Frequency	Adverse drug reactions
Nervous system disorders		Uncommon	headache dysgeusia
Eye disorders		Common	ocular discomfort dry eye blurred vision eye pruritus ocular hyperemia eyelid margin crusting
Respiratory, thoracic and mediastinal disorders		Uncommon	nasal dryness
Gastrointestinal disorders		Rare	dry mouth

Adverse drug reactions from spontaneous reports and literature cases (frequency not known)

The following adverse drug reactions have been derived from post-marketing experience with Pataday via spontaneous case reports and literature cases. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency which is therefore categorized as not known. Adverse drug reactions are listed

according to system organ classes in MedDRA. Within each system organ class, ADRs are presented in order of decreasing seriousness.

Table 0-2 Adverse drug reactions from spontaneous reports and literature (frequency not known)

System Organ Classification	Adverse drug reactions
Immune system disorders	hypersensitivity
Eye disorders	lacrimation increased
Gastrointestinal disorders	nausea

8 Interactions

No clinically relevant interactions have been described.

9 Pregnancy, lactation, females and males of reproductive potential

9.1 Pregnancy

Risk summary

There is a limited amount of data from the use of olopatadine in pregnant women. Studies in rats and rabbits in which olopatadine was orally administered did not show any embryo fetal toxicity up to 2480-times the maximum recommended ocular human dose (MROHD) (one drop of 0.7 % olopatadine ophthalmic solution in each eye, based on body surface area (BSA)). Reduction in the fetal weight was not observed in rats up to 25 times the MROHD, based on BSA.

No effects during pregnancy are anticipated since systemic exposure to olopatadine is negligible by the topical ocular route. However, the possibility of harm to the fetus cannot be ruled out.

Data

Animal data

In an embryo-fetal development (EFD) study in rats, olopatadine (60, 200 and 600 mg/kg/day) was administered orally throughout the period of organogenesis. Mydriasis, hyperaemia and congestion of the ocular fundus, abnormal respiratory sounds were observed in treated dams at high dose levels and the maternal no-effect dose level was 60 mg/kg/day (corresponding to 746-times the MROHD, based on BSA). In offspring, decrease in body weight of live fetuses and decrease in number of ossification were observed at 600 mg/kg/day (corresponding to 7460-times the MROHD, based on BSA). At 60 mg/kg/day, cleft palate was observed in 2 fetuses but not at higher doses. No dose related abnormalities were observed in external, skeletal and visceral examination and hence the no effect dose for offspring was 200 mg/kg/day (corresponding to 2480-times the MROHD, based on BSA).

In a rabbit EFD study, olopatadine (25, 100 and 400 mg/kg/day) was administered orally during the period of organogenesis. Abnormal respiration and lacrimation was seen at the 400 mg/kg/day dose and the maternal no effect dose level was 100 mg/kg/day (corresponding to 2480-times the MROHD, based on BSA). No effects on the fetuses were observed and hence

the no effect dose for offspring was 400 mg/kg/day (corresponding to 9950-times the MROHD, based on BSA).

In a peri-/postnatal toxicity study, rats received oral doses of olopatadine up to 600 mg/kg/day during late gestation and throughout lactation. Maternal toxicity was observed at 600 mg/kg/day. Olopatadine produced decreased neonatal survival at 60 mg/kg/day and reduced body weight gain in offspring at 4 mg/kg/day (50-times the MROHD, based on BSA) which is attributed to milk as demonstrated in a cross-fostered study (see section 9.2 Lactation - Animal Data).

9.2 Lactation

Risk summary

It is not known if olopatadine is transferred into human milk after administration of Pataday. There are no data on the effects of olopatadine on the breastfed child or on milk production. Based upon the low level of olopatadine present in human plasma following topical ocular administration, the concentration of olopatadine potentially present in breast milk is expected to be negligible. However, as there is no data available on the concentration of olopatadine/metabolites in human milk following topical ocular administration, a risk to the breast-feeding child cannot be excluded.

Olopatadine is transferred into the milk of lactating rats after oral administration and was associated with fetal toxicity (see Animal Data).

Patients should be informed that antihistamines may affect the milk production of a nursing mother. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Pataday and any potential adverse effects on the breast-fed child from Pataday.

Data

Animal Data

In a cross-fostered study in which pups of untreated dams were nursed by olopatadine (60 mg/kg/day) treated dams, the body weight gain of pups was suppressed confirming that the effect of olopatadine was through milk.

Oral administration of 1 mg/kg radiolabelled olopatadine in rats demonstrated that olopatadine and/or its metabolites were significantly transferred into milk with milk:plasma ratio ($AUC_{0-\infty}$) of 1.5. Maximal levels of radioactivity in the milk was determined at around 1 hour post-dose, with an elimination half-life of 28.3 hours.

9.3 Females and males of reproductive potential

Studies have not been performed to evaluate the effect of administration of olopatadine on human fertility. Effects in non-clinical fertility studies in male and female animals were observed only at dosages considered sufficiently in excess of the maximum human exposure indicating little relevance to clinical use.

No effects on human fertility are anticipated since systemic exposure to olopatadine is negligible by the topical ocular route.

Olopatadine can be used by women of childbearing potential.

10 Overdosage

Due to the characteristics of this preparation, no toxic effects are to be expected with an ocular overdose of this product, nor in the event of accidental ingestion of the contents of one bottle.

11 Clinical pharmacology

Pharmacotherapeutic group, ATC

Pharmacotherapeutic group: ophthalmologicals; decongestant and antiallergics; other antiallergics.

ATC code: S01GX09

Pharmacodynamics (PD)

Olopatadine is a potent selective antiallergic/antihistaminic agent that exerts its effects through multiple distinct mechanisms of actions. It antagonizes histamine (the primary mediator of allergic response in humans) and prevents histamine induced inflammatory cytokine production by human conjunctival epithelial cells. Data from in vitro studies suggest that it may act on human conjunctival mast cells to inhibit the release of pro-inflammatory mediators. In patients with patent nasolacrimal ducts, topical ocular administration of olopatadine eye drops, solution was suggested to reduce the nasal signs and symptoms that frequently accompany seasonal allergic conjunctivitis. It does not produce a clinically significant change in pupil diameter.

Pharmacokinetics (PK)

Absorption

Olopatadine was absorbed into the eye and reached maximal levels (C_{max}) within 30 minutes to 2 hours (T_{max}) in ocular tissues following bilateral single topical ocular instillation of 1 drop of increasing dose strengths of olopatadine (0.15%, 0.2% and 0.7%) in Male New Zealand White (NZW) Rabbits. Plasma levels of olopatadine were low ($C_{max} < 20$ ng/mL) following bilateral topical ocular administration of 0.15%/0.2%/0.7% olopatadine ophthalmic solution to rabbits.

In the humans, plasma levels following topical ocular administration -and oral administration are shown in Table 11-1. Compared with the oral administration exposure on Day 12, the mean exposure estimates show olopatadine C_{max} (1.64 ng/mL) and AUC_{0-12} (9.68 ng*h/mL) after multiple 0.77% topical ocular doses was 184-fold and 102-fold lower than the C_{max} (302 ng/mL) and AUC_{0-12} (987 ng*h/mL) after multiple 20 mg oral doses of olopatadine. These data indicate that topical ocular doses of 0.77% olopatadine hydrochloride ophthalmic solution resulted in a systemic exposure that is much lower than that after oral doses of 20 mg olopatadine hydrochloride.

Table 11-1 Comparison of olopatadine plasma concentration after topical ocular dosing and oral dosing

Route of administration	Dosage	C_{max} (ng/mL) Mean $\pm$ SD	AUC (ng*hr/mL) Mean $\pm$ SD
-------------------------	--------	------------------------------------	---------------------------------

Topical ocular	1 drop of 0.77% in both eyes once daily, 6.5 days	1.64 ± 0.889	9.68 ± 4.42
	2 drops of 0.1% in both eyes, 4 times-daily, 4 days	0.565 ± 0.463	1.95 ± 1.28 ^{*1}
	2 drops of 0.15% in both eyes, twice-daily, 14 days	0.76 ± 0.31	- ^{*2}
	2 drops of 0.2% in both eyes, twice-daily, 7 days	0.736 ± 0.327	3.63 ± 1.70 ^{*3}
Oral	20 mg tablet, twice-daily, 13.5 days	302 ± 53	987 ± 146 ^{*3}

^{*1}: AUC₀₋₆ ^{*2}: Not calculated because of insufficiency of samples ^{*3}: AUC₀₋₁₂ mean estimates from Day 12

Distribution

Studies in rabbits show ocular tissues associated with the site of dosing i.e., conjunctiva and cornea, had the highest concentrations of olopatadine after bilateral single topical ocular instillation of 1 drop of increasing dose strengths of olopatadine (0.15%, 0.2% and 0.7%) in Male New Zealand White (NZW) Rabbits. Olopatadine concentrations in aqueous humor, choroids, ICB and lens increases with increasing concentrations of olopatadine. Studies conducted in pigmented Dutch belted rabbits indicated a low degree of binding to melanin pigmented tissues.

Biotransformation/metabolism

Studies have not been conducted to investigate the metabolism of olopatadine in ocular tissues. The major metabolites of olopatadine following oral administration in humans are N-desmethyl Olopatadine (M1) and Olopatadine N-oxide (M3). N-desmethyl Olopatadine (M1) is almost exclusively demethylated by the cytochrome P-450 isozyme 3A4 (CYP3A4). Olopatadine was not an inhibitor of cytochrome P-450 isozymes and therefore drug-drug interactions due to metabolic interactions were not expected.

In the humans after topical ocular administration, N-desmethyl metabolite of Olopatadine (M1) was not quantifiable (≤0.050 ng/mL) in plasma sample in all subjects.

Elimination

Studies have not been conducted to investigate the excretion of olopatadine in the urine or feces after topical ocular instillation. In rats after 14C oral administration, olopatadine was rapidly eliminated from the body primarily by urinary excretion and biotransformation (metabolism). In humans, urinary excretion of unchanged drug was the major route of elimination.

Studies conducted to investigate the elimination of olopatadine in rabbits showed concentrations of olopatadine in various ocular tissues (aqueous humor, choroid, conjunctiva, cornea, and ICB) over the dose strengths (0.1 to 0.7% ophthalmic solution) declined with a half-life of less than 4.65 hours.

In humans, the systemic plasma half-life was less than 3 hours.

Linearity/non-linearity

In a single dose study, olopatadine showed a dose proportional increase in exposure (C_{max} and AUC) in ocular tissues after topical ocular instillation.

12 Clinical studies

Pataday is a well-established product.

13 Non-clinical safety data

Non-clinical data reveal no special hazard for humans treated with olopatadine hydrochloride eye drops, solution at concentrations up to and including 0.7% based on conventional studies of single dose toxicity, repeated dose toxicity, genotoxicity, carcinogenic potential and in ocular irritation studies.

For information on embryo-fetal, peri- and post-natal toxicity, see section 9 Pregnancy, lactation, females and males of reproductive potential.

14 Pharmaceutical information

Incompatibilities

None known.

Special precautions for storage

Do not store above 30°C. Discard 28 days after opening.

Instructions for use and handling

No special requirements.

Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

15 Product Registration Holder

Novartis Corporation (Malaysia) Sdn. Bhd.
Level 18, Imazium,
No. 8, Jalan SS21/37, Damansara Uptown,
47400 Petaling Jaya, Selangor

Tracking Number

NA

Information issued

17 Jul 2020

Date of revision

14 Jan 2025