

FUSIGEN*
(Sodium Fusidate Ointment BP)

Name of product: FUSIGEN OINTMENT

Name and Strength of Active Ingredient:

Each g contains: Sodium Fusidate BP 20mg in a neutral ointment base

Dosage Form: Topical Ointment

Product Description: Fusigen Ointment is a white to pale yellow, soft and smooth greasy ointment

Pharmacodynamics:

FUSIGEN is a topical antibacterial agent.

Fusidic acid and its salts show strong surface activity and are also fat soluble. These properties facilitate its diffusion between aqueous and lipid phases in complex media. This may account in part for the tissue penetrating properties exhibited by the antibiotic. (FUSIGEN has been shown experimentally to penetrate skin as effectively as the corticosteroids).

Concentrations of 0.03 – 0.12 mcg fusidic acid per ml inhibit nearly all strains of *Staphylococcus aureus*. Topical application of fusidic acid is also effective streptococci, corynebacteria, nisseria and certain clostridia.

Pharmacokinetics:

Fusidic acid is well absorbed from the gastrointestinal tract, highly bound to plasma proteins (97% to 99.8%), distributed to many tissues, and metabolized in the liver to inactive metabolites. The drug has a serum half-life of 5 to 6 hours and is eliminated almost exclusively in bile

Indications:

Indicated in the treatment of skin infections caused by Staphylococci

Recommended Dosage:

Adults and children:

Uncovered lesions – a small quantity should be applied to the affected area twice daily until a satisfactory response is obtained. A single treatment course should not normally exceed 7 days

Covered lesions – less frequent applications may be adequate.

Route of administration:

Topical application

Contraindications:

Infection caused by non-susceptible organisms, in particular *Pseudomonas aeruginosa*.

Hypersensitivity to fusidic acid and its salts. As with other topical corticosteroid preparations, FUSIGEN OINTMENT is contraindicated in primary bacterial, viral and fungal skin infections, skin manifestations in relation to tuberculosis or syphilis, perioral dermatitis and rosacea.

Warnings & Precautions:

Warnings

Bacterial resistance has been reported to occur with the use of fusidic acid applied topically. As with all topical antibiotics, extended or recurrent application may increase the risk of contact sensation and the development of antibiotic resistance.

The sodium salt of fusidic acid has been shown to cause conjunctival irritation. The ointment should not be used near the eye.

Precautions

Caution should be exercised when using FUSIGEN Ointment near the eye.

Interactions with other medicaments: Not applicable

Pregnancy and Lactation:

There is inadequate evidence of safety in human pregnancy. Animal studies and many years of clinical experience have suggested that fusidic acid is devoid of teratogenic effect.

Safety in nursing mothers has not been established. When fusidic acid (as the sodium salt) has been given systemically, levels have been detected in breast milk, but with topical use the possible amount of drug presents is unlikely to affect the infant.

Side Effects:

Hypersensitivity reactions to the active ingredient in the form of skin rashes; mild stinging and irritation on application have been reported rarely.

Effects on ability to drive and use machinery: Not applicable

Symptoms and Treatment of Overdose: Not applicable

Storage conditions: Store at a temperature not exceeding 30°C.

Shelf life:

The product is stable for a period of 30 months from the date of manufacture.

Available pack sizes: 3gm, 10gm & 15gm

Registration no.: MAL08061611AZ

Mfg. Lic. No. 52

Product Registration Holder:

Komedic Sdn Bhd, 4, Jalan PJS 11/14, Bandar Sunway, 46150 Petaling Jaya, Selangor

Manufactured in India by:

Geno Pharmaceuticals Pvt. Ltd., Tivim Industrial Estate, Karaswada, Mapusa, Goa 403526

Date of Revision: March 2023