



GoldLife® Jointril 500

Description : Red opaque cap/Cream opaque body, size 0 capsule, filled with white to light yellowish powder

Content : Each capsule contains :
Crystalline Glucosamine Sulphate Sodium 628mg
(equivalent to Glucosamine Sulphate 500mg)

Indications : As adjuvant therapy for osteoarthritis.

Pharmacology:

Glucosamine is an aminomonosaccharide, particularly in the management of all forms of degenerative osteoarticular diseases e.g. arthrosis and osteoarthritis, both subacute and chronic, with disappearance or reduction of articular pains, improvement of articular functions and inhibition or regression of the degeneration process. Glucosamine possesses arthrotrophic metabolic activity by stimulating the anabolic metabolism of osteocartilaginous tissues, through the stimulation of the biosynthesis of the mucopolysaccharides (which are essential components of the cartilage substances) and of the bone mesenchymal tissues.

Lubricant activity through improvement of synovial fluid viscosity by increasing synovial fluid production. After oral administration, only 26% is absorbed, with onset of action about 2-3 weeks for osteoarthritis. Bioavailability is low due to first pass hepatic metabolism. Following oral administration, glucosamine is extensively metabolised in the liver. It is rapidly desulfated and metabolised into smaller molecules, and ultimately to carbon dioxide, water and urea. Glucosamine is not protein-bound, but rather incorporates into the plasma proteins (primarily globulin). Rapid diffusion into most tissues also occurs, including articular tissues. Urinary excretion accounts for 10% of an oral dose, about 11% is eliminated via the faeces, presumably as an unabsorbed drug. The elimination half-life of glucosamine which is incorporated into the plasma proteins is about 68 hours after oral doses. Part of a dose of glucosamine sulphate is eliminated as carbon dioxide via expired air in the lungs.

Adverse effects:

Cardiovascular: Peripheral oedema, tachycardia were reported in a few patients following a larger clinical trial investigating oral administration osteoarthritis. Causal relationship has not been established.

Central nervous system: Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

Gastrointestinal: Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy of osteoarthritis.

Skin: Skin reactions such as erythema and pruritis have been rare complications.

Precautions: Glucosamine sulphate is a causal therapy and the therapeutic effect is only seen after 1week. Therefore, it is advisable to include an antiinflammatory drug (in case of severe pains) during the first days of treatment with glucosamine. This product is derived from seafood.

Contraindications: It is contraindicated in patients hypersensitive to glucosamine. Contraindicated in patients allergic to shellfish.

Interaction with other medication : The use of Glucosamine with oral hypoglycemic agent (metformin, tolbutamide, rosiglitazone) may reduce oral hypoglycemic agent effectiveness.

Pregnancy and Lactation : Avoid usage in children, pregnant women and nursing mothers.

Dosage and Administration:

Light to moderate symptoms - 1 capsule twice daily before meals for at least 6 weeks.

Severe symptoms - 1 capsule three times daily before meals.

Capsule should be taken 15 minutes before each meal. Treatment should be repeated every other 6 months or less according to medical prescription.

Storage: Store in a dry place at temperature below 30°C. Protect from light. Keep out of reach of children.

Pack size: Alu-PVC Blister pack - 50's, 60's, 90's
HDPE Bottle pack - 30's, 60's, 90's

Shelf life: 2 years

Contract Manufactured for Medipharma Industries Sdn Bhd
Intan Business Centre 2, 3rd Floor, B-3-9, Lebuhr Medan Ipoh
Bandar Baru Medan Ipoh, 31400 Ipoh, Perak, Malaysia
by Prime Pharmaceutical Sdn Bhd