BiO-LiFE®

Malaysia for submission

Revision Date : 19/09/2017
Revision History :
9/03/2017 - Update Address and Phone No (for submission)
19/09/2017 - add in Mega , update new phone no

Name of Product / Pack size: Glucosamine 500mg with Chondroitin Sulphate 400mg / 90 Tablets

Note: This Label is applicable for all pack sizes

MEGA We care

BiO-LiFE®

Glucosamine
500mg with
Chondroitin
Sulphate 400mg

As an adjunctive therapy in **osteoarthritis**

90
Tablets

One tablet contains :

Glucosamine hydrochloride	500mg
Chondroitin sulphate sodium	445mg
Equiv to chondroitin sulphate 400mg	

No added sugars, yeast, milk derivatives, wheat or corn starch, gluten, preservatives & artificial flavours.

Indication : As an adjunctive therapy in osteoarthritis.

Directions for use :
Light to moderate : 1 tablet twice daily, 15 minutes before meals for at least 6 weeks (or according to medical prescription)
Severe osteoarthritis symptoms:
Initial therapy : 1 tablet 3 times daily, 15 minutes before meals during a period of at least 8 weeks
Follow-up therapy : 1 tablet twice daily, 15 minutes before meals during a period of at least 8 weeks.
Please refer detail dosage information inside flap.

Osteoarthritis, also called degenerative joint disease is caused by the breakdown of cartilage which is the connective tissue that cushions the ends of bones within the joint. As the cartilage breaks down, it loses its ability to cushion the bones resulting in pain.

Glucosamine & Chondroitin sulphate are both substances used in the treatment of osteoarthritis. They are naturally found in and around the cells of cartilage.

Glucosamine:
An amino sugar that the body produces and distributes in cartilage and other connective tissue for building the cartilage.

Chondroitin sulphate:
A complex carbohydrate that helps cartilage retain water.

Store below 30°C in a dry place, away from direct sunlight.
Keep out of reach of children. (Jauhi dari kanak-kanak.)

Ⓑ
Mfg Date :
Exp Date :
Reg No : MAL08111784XC

MS 2424:2012
1 019-07/2011

Marketing Authorization Holder
BiO-LiFE Marketing Sdn Bhd (167808-V)
B-09-01, The Ascent, Paradigm.
No.1, Jalan SS7/26A, Kelana Jaya,
47301 Petaling Jaya, Selangor, Malaysia.
For information on health, please call us
at 03-7499 7999, fax at 03-7882 6388 or
e-mail us at health@biolife.com.my

Manufactured by
Unison Nutraceuticals Sdn. Bhd (298703-W),
No 13, Jalan TTU 52, Taman Tasik Utama,
75450 Ayer Keroh, Melaka, Malaysia
for BiO-LiFE Marketing Sdn Bhd

Chondroitin sulphate sodium derived from bovine.
Glucosamine hydrochloride derived from seafood.

BiO-LiFE® is a registered trademark of BiO-LiFE Marketing Sdn. Bhd.

OPEN HERE

Description: Each tablet contains Glucosamine Hydrochloride 500mg and Chondroitin Sulphate Sodium 445mg in 16.6mm oval shaped white film coated tablets.

Route of Administration: Oral

Indication: As an adjunctive therapy in osteoarthritis

Directions for use :
Light to moderate : 1 tablet twice daily, 15 minutes before meals for at least 6 weeks (or according to medical prescription)
Severe osteoarthritis symptoms:
Initial therapy : 1 tablet 3 times daily, 15 minutes before meals during a period of at least 8 weeks.

Follow up therapy: Maintenance therapy should be followed for 3-4 months (or according to medical prescription) by administration of 1 tablet taken 2 times daily, 15 minutes before meals. The treatment should be repeated every other 6 months or less according to medical prescription.

Mode of action: The glucosamine and chondroitin sulphate that are taken up to the joints stimulates biosynthesis of mucopolysaccharide which is an essential component of the cartilage. They are believed to play a role in the promotion and maintenance of the structure and function of cartilage. Their action includes pain relief of osteoarthritic joint and anti-inflammatory activity.

Pharmacokinetics: Upon consumption of the BiO-LiFE Glucosamine 500mg and Chondroitin Sulphate 400mg, the tablet disintegrates in the stomach to release glucosamine & chondroitin sulphate. About 90% of glucosamine administered orally as a glucosamine salt gets absorbed from the small intestine and from there it is transported via portal circulation to the liver. Some amount of the ingested glucosamine is catabolized in liver while some is taken up in the joints. Exact percentages that are catabolized or delivered to the joints are not known. Absorption of chondroitin sulphate occurs both from the stomach and small intestine. There are indications from studies which says after absorption chondroitin sulphate does enter the joint space.

Contraindications: Contraindicated to those who are hypersensitive to glucosamine or other ingredient in the product or allergic to shellfish.

Warning and precautions: This product contains glucosamine. Those with type 2 diabetics and those who have problems with glucose tolerance should have their blood sugars carefully monitored if they use glucosamine. Please include anti - inflammatory drug (in case of severe pain) during first day of treatment. Glucosamine and Chondroitin are not meant to replace NSAIDS in cases of intense pain as this is a causal therapy.

Pregnancy and lactation: This product contains chondroitin sulphate which is not recommended for children, pregnant women and nursing mothers.

Interactions with other medicaments:
Diabetics on blood sugar control drugs should take this supplement under medical supervision. There is a possibility that chondroitin sulphate may have antithrombotic activity. Those taking warfarin and those with haemophilia should exercise caution in its use. Chondroitin should not be used concomitantly with chitosan.

Symptoms and treatment of overdose:
In the event of an acute overdose, the patient should be observed closely and symptomatic treatment should be instituted.

Side effects: Mild gastrointestinal complaints such as heartburn, epigastric distress, nausea and diarrhoea have been reported. No sulfa-allergic reactions or other allergic reactions have yet been reported.

Cardiovascular: Peripheral oedema, tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

Central Nervous System: Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

Gastrointestinal: Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine.

Skin: Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

Dosage Form: Tablet

Packing Sizes: 30s, 90s

Marketing Authorization Holder: BiO-LiFE Marketing Sdn Bhd (167808-V),
B-09-01, The Ascent, Paradigm. No.1, Jalan SS7/26A, Kelana Jaya, 47301 Petaling Jaya, Selangor, Malaysia.

Manufactured by: Unison Nutraceuticals Sdn. Bhd (298703-W)
No 13, Jalan TTU 52, Taman Tasik Utama, 75450 Ayer Keroh, Melaka, Malaysia

Revision Date: 19th September 2017