

Front

60 mm

60 mm

240 mm

For Hospital or Medical practitioner or Laboratory

CONTRAPAQUE 300

CONTRAPAQUE 350

(Iohexol Injection USP)

Name and strength of Active ingredients for Contrapaque 300 mg:

Each ml contains:

Iohexol USP

647 mg

Equivalent to Iodine

300 mg

Water for injection BP

q.s

Name and strength of Active ingredients for Contrapaque 350mg:

Each ml contains:

Iohexol USP

755 mg

Equivalent to Iodine

350 mg

Water for injection BP

q.s.

Description:

Clear, colourless to pale yellow slightly viscous liquid, free from foreign matter.

Pharmacodynamics:

Iohexol is a "second generation nonionic radio contrast agent. It is available for intravenous, intra-arterial and intra-thecal administration in diagnostic radiology. In comparison with conventional (ionic) radiocntrast media (eg, metrizoate, iothalamte and diatrizoate) iohexol has a relatively low osmolality. This lower osmolality has been associated with a lower incidence of side effects such as pain, heat sensations, adverse hemodynamic changes, EKG changes, endothelial damage and damage to erythrocytes.

Pharmacokinetics:

Absorption

•

Following intrathecal injection, iohexol is rapidly absorbed into the bloodstream from CSF. Peak plasma concentrations occur between 2 and 6 hours after injection.

Distribution

•

Iohexol has low affinity for serum or plasma proteins and is poorly bound to serum albumin.

•

After intravascular injection iohexol is distributed in the extra cellular fluid compartment, with no significant deposition. Iohexol does not cross the blood- brain barrier to any significant extent following intravascular administration. Iohexol probably crosses the placental barrier in humans by simple diffusion. The volume of distribution is reported to be 0.577 L/kg (0.35 to 0.849 L/kg) after lumbar intrathecal injection.

•

After intravenous injection, the volume of distribution has been reported to be 0.165 L/kg (0.108 to 0.219 L/kg)

Metabolism

•

Iohexol undergoes minimal metabolism, de-iodination, or bio-transformation. No metabolites of iohexol have been detected in urine, feces or plasma.

Excretion

•

Following intrathecal or intravenous administration, iohexol is excreted primarily through the kidneys, almost entirely via glomerular filtration. In patients with severe renal insufficiency or failure, compensatory biliary excretion of the drug is anticipated to occur, with a slow clearance into the bile. It is not known to what extent iohexol is excreted in human milk.

Half-Life

The elimination half-life to iohexol, after intravenous administration, is approximately 2 hours. After intrathecal administration, terminal elimination half-life is 4.5 hours.

Indication

X-ray contrast medium for use in adults and children for cardioangiography, arteriography, urography, phlebography and CT-enhancement. Lumbar, thoracic, cervical myelography and computed tomography of the basal cisterns, following subarachnoid injection. Arthrography, endoscopic retrograde pancreatography, (ERP), endoscopic retrograde cholangiopancreatography (ERCP), herniography, hysterosalpingography, sialography and studies of the gastrointestinal tract.

Recommended Dose

The Dosage vary depending on the type of examination, age, weight, cardiac output and general condition of the patient and the technique used. For intravenous, intra-arterial and intrathecal use, and use in body cavities.

Guidelines for Intravenous use

Indication	Concentration	Volume	Comments
Urography			
<u>adults:</u>	300 mg I/ml	40 - 80 ml	80 ml may be exceeded in selected cases
	or 350 mg I/ml	40 - 80 ml	
<u>children < 7 kg</u>	240 mg I/ml	4 ml/kg	
	or 300 mg I/ml	3 ml/kg	
<u>children > 7 kg</u>	240 mg I/ml	3 ml/kg	
	or 300 mg I/ml	2 ml/kg (max 40 ml)	
Phlebography (leg)	200 mg I/ml	20 - 100 ml/leg	
	or 240 mg I/ml		
	or 300 mg I/ml		
Digital subtraction angiography	300 mg I/ml	20 - 60 ml/inj.	
	or 350 mg I/ml	20 - 60 ml/inj.	
CT-enhancement	140 mg I/ml	100 - 400 ml	
<u>adults:</u>	or 200 mg I/ml	100 - 300 ml	Total amount of iodine usually 30 - 60 g
	or 240 mg I/ml	100 - 250 ml	
	or 300 mg I/ml	100 - 200 ml	
	or 350 mg I/ml	100 - 150 ml	
<u>Children:</u>	240 mg I/ml	2-3 ml/kgbw up to 40 ml	In a few cases up to 100 ml may be given
	or 300 mg I/ml	1-3 ml/kgbw up to 40 ml	

Guidelines for Intrathecal use

Indication	Concentration	Volume	Comments
Lumbar and thoracic myelography	180 mg I/ml	10 - 15 ml	
(lumbar injection)	or 200 mg I/ml	10 - 15 ml	
	or 240 mg I/ml	8 - 12 ml	
Cervical myelography (lumbar injection)	240 mg I/ml	10 - 12 ml	
	or 300 mg I/ml	7 - 10 ml	
Cervical myelography (lateral cervical injection)	240 mg I/ml	6 - 10 ml	
	or 300 mg I/ml	6 - 8 ml	
CT cisternography (lumbar injection)	180 mg I/ml	5 - 15 ml	
	or 200 mg I/ml	5 - 15 ml	
	or 240 mg I/ml	4 - 12 ml	
Paediatric myelography			
<2 years	180 mg I/ml	2 - 6 ml	
2-6 years	180 mg I/ml	4 - 8 ml	
>6 years	180 mg I/ml	6 - 12 ml	
To minimize possible adverse reactions a total dose of 3 g iodine should not be exceeded.			

Guidelines for Intra-arterial use

Indication	Concentration	Volume	Comments
Arteriographies			
arch aortography	300 mg I/ml	30 - 40 ml/inj	Volume pr. injection depends on the site of injection
selective cerebral	300 mg I/ml	5 - 10 ml/inj	
aortography	350 mg I/ml	40 - 60 ml/inj	
femoral	300 mg I/ml	30 - 50 ml/inj	
	or 350 mg I/ml		
various	300 mg I/ml	depending on type of examination	
Cardioangiography			
<u>adults:</u>			
left ventricle and aortic root inj.	350 mg I/ml	30 - 60 ml/inj.	
<u>selective coronary arteriography</u>	350 mg I/ml	4 - 8 ml/inj.	
<u>children:</u>		depending on age, weight and pathology	
	300 mg I/ml	(max 8 ml/kg)	
	or 350 mg I/ml		
Digital subtraction angiography	140 mg I/ml	1 - 15 ml/inj.	depending on site of inj. occasionally large volumes - up to 30 ml - may be used
	or 200 mg I/ml	1 - 15 ml/inj.	
	or 240 mg I/ml	1 - 15 ml/inj.	
	or 300 mg I/ml	1 - 15 ml/inj.	

Guidelines for Body cavities

Indication	Concentration	Volume	Comments	
Arthrography	200 mg I/ml	5 - 20 ml		
	or 240 mg I/ml	5 - 20 ml		
	or 300 mg I/ml	5 - 15 ml		
	or 350 mg I/ml	5 - 10 ml		
ERP/ERCP	240 mg I/ml	20 - 50 ml		
Herniography	240 mg I/ml	50 ml	The dosage varies with the size of the hernia	
Hysterosal-pingography	240 mg I/ml	15 - 50 ml		
	or 300 mg I/ml	15 - 25 ml		
Sialography	240 mg I/ml	0.5 - 2 ml		
	or 300 mg I/ml	0.5 - 2 ml		
Gastrointestinal studies				
Oral use				
<u>Adults:</u>	180 mg I/ml	individual		
	or 200 mg I/ml	individual		
	or 350 mg I/ml	individual		
<u>Children:</u>				
- esophagus	300 mg I/ml	2-4 ml/kgbw	Max. dose 50 ml	
	or 350 mg I/ml	2-4 ml/kgbw	Max. dose 50 ml	
- ventricle/follow trough	140 mg I/ml	4-5 ml/kgbw		
<u>Prematures:</u>	350 mg I/ml	2-4 ml/kgbw		
Rectal use	140 mg I/ml	5-10 ml/kgbw	Example: Contrapaque 240, 300 or 350 with top-water 1:1 or 1:2	
- children:	or dilute with tapwater to 100-150 mg I/ml	5-10 ml/kgbw		
CT-enhancement				
Oral use				
- adults:	Dilute with tapwater to ~6mgI/ml	800 -2000 ml of the diluted solution over a period of time	Example: Contrapaque 300 or 350 with top-Water 1:50	
- children:	Dilute with tapwater to ~6mgI/ml	15-20 ml/kgbw of the diluted solution		
Rectal use	Dilute with tapwater	individual		
- children:	to ~6mgI/ml			

Route of administration

For intravenous, intra-arterial and intrathecal use, and use in body cavities.

Contraindication

•

Iohexol should not be administer to patients with known hypersensitivity to Iohexol.

•

Iohexol should not be administered with intrathecal corticosteroids.

•

Immediate repeat myelography with iohexol is contraindicated because of the possibility of overdosage.

•

Myelography is contraindicated in the presence of significant local or systemic infection where bacteremia is likely.

220 mm

SAP Code:142792	Country :Malaysia	Mfg. Location :Panoli IV17	Product Name : Contrapaque 300-350
Packaging Material Leaflet	Existing / Reference Art Party's Word File	Version No.: 1	Date : 03-11-2025
Dimension : 220 (L) x 240 mm (B) (± 1 mm tolerance)		Size After Folding :55 x 60 mm (± 1 mm tolerance)	Core Dia : NA
Reel Dia NA	Varnish / Lamination :NA	Roll Unwind Direction :NA	Print Repeat Length :NA
Specification : ITC Fine paper		Pantone : Black	Folding pattern : 2 Vertical + 2 Horizontal Folds
Grammage: 40 ± 5% gm/m ²			
Thickness NA			
Other Test :			
Reason for Change New Artwork		Unique J. B. CHEMICALS & PHARMACEUTICALS LTD.	
File Path : X:\Artworks\Mahesh\Malaysia\Contrapaque V2\Open\ 142792 Contrapaque 300-350 Leaflet.ind			

Artist : VIJAY

This document is equivalent to a digitally certified copy & digital signature is not required.

