

Revision Date : 06-01-2025
Revision Date : 23-10-2023
Revision Date : 06/08/2024

Artwork is in same size
Size: 90 mm x 200 mm

For Hospital or Medical practitioner or Laboratory

MUCOPAIN Gel

(Benzocaine USP 20% w/w)

Name and strength of Active ingredient(s):

Benzocaine USP 20% w/w

Description:

A slight yellow clear gel with whitish fluorescence and characteristic odour

Pharmacodynamics:

Benzocaine is a surface anesthetic of ester type. It has found frequent use to treat pain arising from various mucosal membranes.

Pharmacokinetics:

Benzocaine is sparingly soluble in water with toxicity about the tenth (1/10) that of cocaine. It is not readily absorbed from mucous membranes. It is an ester, which on hydrolysis produces p-aminobenzoic acid.

Indication

To be applied topically to oral tissues for temporary relief of pain associated with dental conditions.

Recommended Dose

Apply small amount evenly to the affected area as directed by the physician.

Route of administration

Topical or external use only

Contraindication

Patients with known hypersensitivity to Benzocaine. Application to severely traumatized mucosa or to areas of sepsis.

Warning and precaution

Localized allergic reactions may occur in patients sensitive to Benzocaine or after prolonged or repeated use. Methemoglobinemia may rarely occur, particularly in infant and children, following application of large amounts e.g. in pharyngeal anesthesia. Topical benzocaine should be used in children below 2 years with caution. It should not be used for prolonged periods.

Interaction with other medicaments

Cross-sensitivity may occur with other local anesthetic of the ester type and systemic medications such as sulphonamides, thiazides and para-amino dyes.

Pregnancy and lactation

Animal reproduction studies have not been conducted with benzocaine. Mucopain Gel should be given to a pregnant woman only if clearly indicated. It is not known whether benzocaine is excreted in human milk. Caution should be exercised when Mucopain Gel is administered to a nursing woman.

Side effects

Contact dermatitis and/or hypersensitivity to benzocaine can cause burning, stinging, pruritus, tenderness, erythema, rash, urticaria and edema. Rarely, benzocaine may induce methemoglobinemia causing respiratory distress and cyanosis.

Symptoms and treatment of overdose

Accidental swallowing of large quantities causes numbness, nausea, vomiting, irritation to eyes. In case of eye and skin contact, flush with large quantities of water. In case of accidental swallowing, to get medical attention immediately. Treatment is primarily supportive. It includes monitoring for development of methemoglobinemia.

Storage condition

Store in a dry place at a temperature not exceeding 30°C, protect from light.

Shelf life

36 months from the date of manufacture

Presentation

7.5/15 g multicoloured laminated tube with white, tightly fitted cap. Text printed in multicolour and two number dark blue eye marks packed in multicoloured carton.

Manufactured By

ICPA Health Products Limited
286/287/288, GIDC, Ankleshwar
Gujarat, 393002, India
Revision date: 06/08/24

Product registration holder:

UNIMED SDN BHD
No 53, Jalan Tembaga SD 5/2B,
Bandar Sri Damansara,
52200 KUALA LUMPUR

13000201