



SODIUM FUSIDATE- 250 TABLET

COMPOSITION:

Each tablet contains
Sodium Fusidate 250mg

PRESENTATION:

Off white, round biconvex tablet with 10.0mm in diameter.

INDICATIONS:

Sodium Fusidate is indicated for the treatment of infections caused by susceptible organisms, especially staphylococci, e.g. osteomyelitis, septicaemia, endocarditis, superinfected cystic fibrosis, pneumonia, skin and soft tissue infections, surgical and traumatic wound infections.

PHARMACOLOGY:

Mechanism of Action:

Sodium Fusidate is an antibacterial with bacteriostatic or bactericidal activity, mainly against Gram-positive bacteria. Sodium Fusidate inhibits bacterial protein synthesis by preventing the translocation of peptide subunits and elongation of peptide chain. It is active against the following bacteria.

- Gram-positive bacteria : staphylococci (eg. *Staph aureus* and *Staph epidermis*), *Nocardia asteroides* & many clostridial strains. Streptococci and enterococci are less susceptible.
- Gram-negative bacteria : *Neisseria spp.*, *Bacteroides fragilis* and some activity against strains of *Mycobacterium tuberculosis*.

Pharmacokinetics:

Sodium Fusidate is well absorbed from the gastrointestinal tract. Most fusidate in the circulation is bound to plasma protein. Thus, is broadly distributed into tissues and body fluids including bone, pus and synovial fluid. Although it penetrates cerebral abscesses, it does not enter cerebrospinal fluid in appreciable amount. It has been found in the fetal circulation and in breast milk. Single 500mg dose is reported to produce mean plasma concentrations of about 30µg per ml within 2 to 4 hour after administration, although there is considerable interindividual variation. Food may delay its absorption and children may have more rapid absorption than adults. Some accumulation occurs with repeated administration or more have been reported following 500mg of Sodium Fusidate three times daily for 4 days. Fusidate is excreted in

the bile, almost entirely as metabolites. About 2% appears unchanged in the faeces. Little is excreted in the urine or removed by haemodialysis.

DOSAGE AND ADMINISTRATION:

For oral administration

Adult: 500mg (2 tablets) three times daily.

Skin and soft tissue infections: 250mg (1 tablet) twice daily. Doses may be doubled in severe cases.

CONTRAINDICATION:

Hypersensitivity to Sodium Fusidate.

WARNING AND PRECAUTION:

Should be used with caution to patients with hepatic impairment, biliary obstruction or biliary disease, or when it is given in combination with other antibiotic, which have similar excretion pathways. Periodic monitoring of hepatic function is recommended in patient with hepatic impairment and in those receiving high or prolonged oral doses. In vitro, Sodium Fusidate competes with bilirubin for binding to albumin. The clinical significance of this finding is uncertain and kernicterus has not been observed in neonates receiving Sodium Fusidate. However, this observation should be borne in mind when the drug is given to premature, jaundiced, acidotic or seriously ill neonates.

Use in Pregnancy and Lactation:

Animal studies and clinical experience recommend that Sodium Fusidate is lacks of teratogenic effects. It passes the placenta and should be avoided during third trimester due to theoretical risk of kernicterus. Concentration of Sodium Fusidate found in breast milk are negligible and its use is not contraindicated in nursing mothers.

SIDE EFFECT:

May cause mild gastrointestinal disturbance such as dyspepsia, nausea, vomiting, epigastric pain, anorexia, which can normally be avoided by giving it with meals. Treatment with Sodium Fusidate has been associated with jaundice and change of liver function; normal liver function is usually restored when treatment is stopped. Allergic reactions are reportedly very rare.

DRUG INTERACTIONS:

A drug that is metabolized by hepatic cytochrome P450 isoenzymes may be suspected.

OVERDOSAGE & TREATMENT:

Treatment should be restricted to symptomatic and supportive measures. Dialysis is of no benefit as the drug is not significantly dialysed.

STORAGE:

Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

PACK QUANTITIES:

Available in blister pack of 2 x 10's.

Further information can be obtained from pharmacist, doctor or the manufacturer.

MAL09020977AZ

Manufacturer and Product Registration Holder



Kotra Pharma (M) Sdn. Bhd.
No. 1, 2&3, Jalan TTC 12, (00000-1)
Cheng Industrial Estate,
75250 Melaka, Malaysia

REGN-260121

Date of revision: 26/07/19