



OXYMETAZOLINE 0.05% NASAL SPRAY

MAL09032007AZ

COMPOSITION:

Oxymetazoline HCl 0.05% w/v in buffered aqueous solution.
Benzalkonium Chloride 0.01% w/v as preservative.

PRESENTATION:

Colourless liquid.

INDICATIONS:

Symptomatic relief of nasal & nasopharyngeal congestion.

PHARMACOLOGY:*Mechanism of Action:*

Oxymetazoline is a direct - acting sympathomimetic agent which has a vasoconstrictor effect on mucosal blood vessels. Sympathomimetic agent mimic the actions produced by the stimulation of postganglionic sympathetic or adrenergic nerves, including vasoconstriction of blood vessels supplying skin and mucous membranes. Topical use of Oxymetazoline has the advantage of selective site of action and more rapid action than do oral agents. It acts within a few minutes and the effects last for up to 12 hours. Nasal congestion is frequently a symptom of conditions such as rhinitis. They act by vasoconstriction and redistributing local blood flow to reduce oedema of the nasal mucosa thus improving ventilation, drainage and nasal stuffiness.

Pharmacokinetics:

Most sympathomimetic agents undergo enzymatic degradation in the gut and first pass metabolism of the liver. They are inactivated by the enzymes catechol-O-methyltransferase (COMT) and monoamine oxidase (MAO). The metabolites are excreted in the urine mainly as their glucuronide and ethereal sulphate conjugates. Oxymetazoline, however, is an imidazoline derivative and is more resistant to the inactivating enzymes of liver and other tissues. It is thus longer action and metabolized at a slower rate than the catecholamines such as ephedrine or phenylephrine. When used in the form of nasal spray it acts with a few minutes and the effects lasts for several hours.

ROUTE FOR ADMINISTRATION:

For intranasal administration only

DIRECTION FOR USE:

Adult & children above 6 years: 2 - 3 sprays into each nostril twice a day, morning and evening. With the head upright, place the spray nozzle into the nostril without completely occluding the nostril. During each administration, the patient should bend his head slightly forward and press the pump spray quickly and firmly to spray the recommended number of sprays in each nostril. Wipe nozzle clean after use.

CONTRAINDICATION:

Hypersensitive to any of the ingredients.

PRECAUTIONS:

Do not exceed the recommended dosage. Prolonged use of nasal decongestant spray may cause rebound congestion and drug-induced rhinitis. Avoids excess use. Use is therefore

restricted to periods of not more than 3 consecutive days. This product should not be used by patients who have heart disease, high blood pressure, thyroid disease, diabetes or difficulty in urination due to enlargement of the prostate gland without careful clinical consideration. The dosage of Axcel Oxymetazoline 0.05% Nasal Spray for children below 6 years of age has not been established. Use of product dispenser by more than one person may spread infection.

Use in Pregnancy and Lactation:

Safe use of Oxymetazoline hydrochloride in pregnancy and lactation has not established. Hence, it is not to be used during pregnancy and lactation unless the potential benefits outweigh the risk.

SIDE EFFECTS:

Oxymetazoline may occasionally cause local stinging or burning, sneezing and dryness of the mouth and throat. Prolonged or excessive use may cause rebound congestion and drug induced rhinitis.

DRUG INTERACTION:

Concomitant use of tricyclic antidepressants, maprotiline or monoamine oxidase inhibitors (MAOI), depending on the amount present, may also interact with this medication.

OVERDOSAGE AND TREATMENT:

Overdosage may cause CNS depression, leading to coma and marked reduction in body temperature, especially in infants. In case of overdosages, the appropriate symptomatic and supportive treatment should be given. Consult a physician for more information on the management of overdose.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

PACK QUANTITIES:

Available in 10ml amber glass bottle with nasal pump spray.

Further information can be obtained from doctor, pharmacist or the manufacturer.

Product Registration Holder & Manufactured By :



Kotra Pharma (M) Sdn. Bhd.

No. 1, 2 & 3, Jalan TTC 12, (90082-V)

Cheng Industrial Estate,

75250 Melaka, Malaysia.

Date of Revision: 01/07/24