

Malaysia version

Title: Vecobal 500mcg Tablet - PI

Date: 31-3-2017

VECOBAL 500MCG TABLET Active Ingredient: Each tablet contains:- Mecobalamin 500mcg Presentations: Boxes of 6's x 20 and 6's x 100 strips. Product Descriptions: Vecobal 500mcg Tablet is a pink coloured, round, uncoated tablet with flat surface on both sides and score on one side. Pharmacology: Mecobalamin is the one and only homologue of Vitamin B12, found in the brain that participates in transmethylation as a Vitamin B12 containing coenzyme. It plays a fundamental role in protein synthesis and stimulates methionine synthesis from homocysteine and helps to restore normal levels of DNA and RNAs in nerve cells. It also enhances the formation of lecithin (another major component of the myelin sheath). Evidence indicates Mecobalamin is utilized more efficiently than Cyanocobalamin to increase levels of one of the coenzyme forms of Vitamin B12. Absorption occurs almost exclusively in the terminal ileum by an active saturable process, but large amounts (>30mcg) may also be absorbed by passive diffusion (a maximum of 1.5mcg may be absorbed from oral doses of 5 – 50mcg). For normal absorption, the vitamin must bind to salivary haptocorrin and then to 'intrinsic factor', a highly specific glycoprotein secreted by the parietal cells of the stomach. Vitamin B12 is stored mainly in the liver. In the blood, it is bound to specific plasma proteins (transcobalamins). Elimination is via the urinary, biliary and faecal routes. Enterohepatic recycling serves to conserve B12. Vitamin B12 appears in breast milk. Indications: Treatment of peripheral neuropathies. Treatment of symptoms due to Vitamin B12 deficiency. Dosage and Administration: Adults: Take 1 tablet (500mcg) 3 times daily. The dose may be adjusted according to age of the patient and severity of symptoms. Symptoms and Treatments of Overdose: No overdose of Mecobalamin has been documented. In the event of such incident, supportive or symptomatic treatment is recommended. Contraindication: Not recommended for children unless advised by a doctor. Discontinued if symptoms of hypersensitivity occur.	 Precautions: Mecobalamin should not be used continuously for months if there is no response at all after initial administration. If the condition does not improve or symptom persists, please consult a doctor. Prolonged use of larger doses of Mecobalamin is not recommended for patients whose occupation requires handling mercury or its compound. Use in Pregnancy and Lactation: Not recommended during pregnancy and lactation. Vitamin B12 is distributed into breast milk. Interaction with Other Medicaments: Absorption of Vitamin B12 from the gastrointestinal tract may be reduced by Neomycin, Aminosalicylic Acid, Histamine H2-Antagonists and Colchicine. Undesirable Effects: Generally, Mecobalamin is well-tolerated. In rare cases, some GI discomforts (such as anorexia, nausea, vomiting, or diarrhoea) and skin rash may occur. Storage: Store below 30°C. This product is unstable in light. Protect from light and moisture. Always keep in its original packaging provided. Keep out of reach of children. Jauhkan daripada kanak-kanak. Shelf Life: 3 years from the date of manufacture. Registration No.: MAL 09032033XC Product owner:  Pharm-D Health Science Sdn. Bhd. 200101019018 (554775-K) No 8A, Jalan 1/137C, Bedford Business Park, Off Jalan Kelang Lama, 58000 Kuala Lumpur, Malaysia Manufactured by: Winwa Medical Sdn. Bhd. 198101009677 (75794-A) 1952, Lorong I.K.S. Bukit Minyak 3, Taman I.K.S. Bukit Minyak, 14000 Bukit Mertajam, Penang, Malaysia. Product registration holder:  Eldon Healthcare Sdn. Bhd. 200901039772 (882915-A) a member of the pharm-d health science group No 84, Jalan I-Park 1/9, Kawasan Perindustrian I-Park, Bandar Indahpura, 81000 Kulai, Johor, Malaysia. T 07-660 0960 E info@eldon.com.my Version No.: X.TPdVB-I.05 Date of Revision: 28 July 2025
---	---

Size: 75mm x 105mm
 Dark Blue (Pantone 2767U)