130 mm

120 mm

PRESCRIBING INFORMATION

For the use of a Registered Medical Practitioner / a Hospital or Laboratory only

Bitvio™

Ointment

(Mometasone Furoate Ointment USP 0.1%)

Composition

Bitvio Ointment :

Mometasone Furoate USP 0.1%w/w

In a ointment base q.s

Dosage Form: Ointment

Product Description

A smooth, uniform, white to off white opaque ointment, free from foreign matter

Pharmacodynamics

Mometasone furoate, a synthetic corticosteroid exhibits its anti-inflammatory, antipruritic and vasoconstrictive properties.

Pharmacokinetics

Systemic absorption following topical application of mometasone furoate 0.1% is minimal (approximately 0.4% of the applied dose), the majority of which is excreted within 72 hours following application.

Indication

Indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid responsive dermatoses.

Recommended Dose

A thin film of the ointment should be applied to the affected skin areas once daily until the lesion heals or for duration of three weeks, whichever is sooner.

Mode Of Administration

Topical

Contraindication

Mometasone Furoate Ointment 0.1% is contraindicated in those patients with a history of hypersensitivity to any of the components in the preparation.

Warning and Precautions

Side effects that have been reported after systemic use of corticosteroids, including adrenal suppression, may also occur with topical corticosteroids, especially in infants and children.

Application of topical corticosteroids over large surface areas of the body or the use of occlusive dressings will be increase their systemic absorption. Paediatrics patients may demonstrate greater susceptibility than mature patients, to topical corticosteroid included HPA axis suppression and Cushing's syndrome. This is because of a larger skin surface area to body weight ratio in paediatric patients. When treating infants and children or in long-term therapy therefore, appropriate precautions should be observed.

Amount of topical corticosteroids used in children should not exceed the least amount necessary for effective therapy. Chronic corticosteroid therapy may impede growth and development of children.

Interactions With Other Medicaments

There are no drug interactions of note with Mometasone Furoate

Pregnancy and Lactation

There is inadequate evidence of safety in human pregnancy. Topical administration of corticosteroids to pregnant animals can cause abnormalities of foetal development including cleft palate and intra-uterine growth retardation. Topical corticosteroids should be used during pregnancy only if the potential benefit justifies potential risk to the fetus. Drugs of this class should not be used on pregnant patients in large amounts or for prolonged periods of time.

It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Mometasone should be administered to nursing mothers only after careful consideration of the benefit/risk relationship.

Side Effects

Local adverse reactions rarely reported include burning, pruritus, tingling/stinging and signs of skin atrophy.

The following local adverse reactions have been reported infrequently with the use of other topical corticosteroids: irritation, hypertrichosis, hypo pigmentation, perioral dermatitis, and allergic contact dermatitis, maceration of the skin, secondary infection, striae and miliaria.

Symptoms And Treatment of Overdose

Symptoms: Excessive, prolonged use of topical corticosteroids can suppress pituitary-adrenal function resulting in secondary adrenal insufficiency.

Treatment: Appropriate symptomatic treatment is indicated. Acute hyper corticoid symptoms are virtually reversible. Treat electrolyte imbalance, if necessary. In cases of chronic toxicity, slow withdrawal of corticosteroids is advised.

Storage Condition

Store at a temperature not exceeding 30°C. Do not freeze.

Shelf Life

36 months from the date of manufacture.

Packaging available

20g aluminium collapasible tube and flower pot type white HDPE cap packed in a unit printed duplex board carton.

Date of Revision : Sept. 2019

Reg. No. : MAL09062111AZ
Mfg. Lic. No. : 28/9/96

Importer & Product registration holder :
Zulat Pharmacy Sdn. Bhd, 23 & 23A,
Jalan Bandar 3,Taman Melawati,
53100 Kuala Lumpur, Malaysia.

torrent PHARMA

Manufactured by :
TORRENT PHARMACEUTICALS LTD.
Plot No. 810, Sector III, Industrial Area, Pithampur,
District Dhar - 454 775 (M.P.) INDIA

xxxxxxxxx-5253

PRODUCT NAME :	Bitvio Ointment	**COUNTRY :** Malaysia	**LOCATION :** Pithampur		**Supersedes A/W No.:** ZB0EXX0-MY		
ITEM / PACK :	Insert	**NO. OF COLORS:** 1	**REMARK :**				
DESIGN STYLE :	-	**PANTONE SHADE NOS.:**	**SUBSTRATE :**				
CODE :	xxxxxxxxx-5253		**Activities**	**Department**	**Name**	**Signature**	**Date**
DIMENSIONS (MM) :	130 L x 120 W mm	Black C	Prepared By	**Pkg.Dev**			
ART WORK SIZE :	S/S		Reviewed By	**Pkg.Dev**			
DATE :	24-09-2019		Approved By	**Quality**			

This colour proof is not colour binding. Follow Pantone shade reference for actual colour matching.